

ONLINE FIRST PUBLICATION

Online first papers have undergone full scientific review and copyediting, but have not been typeset or proofread. To cite this article, use the DOIs number provided. Mandatory typesetting and proofreading will commence with regular print and online publication of the online first papers of the *SMJ*.

Lateral versus central tendon-splitting approach to insertional Achilles tendinopathy: a retrospective study

Zhan Xia¹, MBBS, Khye Soon Andy Yew¹, BEng, PhD, Ting Karen Zhang², BSc,
Inderjeet Singh Rikhranj¹, FRCS, FAMS

¹Department of Orthopaedic Surgery, ²Orthopaedic Diagnostic Centre, Singapore General Hospital, Singapore

Correspondence: Dr Zhan Xia, Senior Resident, Department of Orthopaedic Surgery, Singapore General Hospital, Outram Road, Singapore 169608. williamxia0402@gmail.com

Singapore Med J 2019, 1–15

<https://doi.org/10.11622/smedj.2019038>

Published ahead of print: 2 May 2019

Online version can be found at
<http://www.smj.org.sg/online-first>

ABSTRACT

Introduction: This study aimed to compare operative outcomes between the lateral approach (LA) and the central approach (CA) to insertional Achilles tendinopathy (IAT).

Methods: We retrospectively reviewed patients who underwent surgical treatment for IAT using the LA or CA. Patients' demographic data, postoperative complications and satisfaction rate were reviewed. Clinical outcomes were prospectively assessed preoperatively, at three months postoperatively and at the last visit, including patients' visual analogue scale (VAS) scores for pain, American Orthopaedic Foot and Ankle Society (AOFAS) Ankle-Hindfoot Scale scores and 36-item Short Form Health Survey (SF-36) scores.

Results: There were 39 patients in the LA group and 32 in the CA group. In each group, average VAS and AOFAS Scale scores improved significantly. SF-36 scores improved in most parameters. No significant difference in baseline score; average AOFAS Scale score at each visit; and mean VAS score preoperatively and at last visit; satisfaction rate; and overall complication rate were observed between the groups. The mean VAS score in the LA group at postoperative three months was significantly lower than that in the CA group (2.7 ± 2.5 vs. 4.4 ± 3.0 ; $p = 0.016$). There were significantly more cases of delayed wound healing in the LA group than in the CA group (2.6% vs. 15.6%; $p = 0.049$).

Conclusion: Both approaches had comparable outcomes for IAT in terms of functional improvement, pain relief, overall enhancement of patients' health condition and overall postoperative complication rate. The LA provided better short-term pain relief and reduced delayed wound healing compared with the CA.

Keywords: Achilles tendon, central tendon-splitting approach, lateral approach, operative procedure, tendinopathy

INTRODUCTION

Insertional Achilles tendinopathy (IAT), a disease at the insertional area of the Achilles tendon into the calcaneum, is characterised by pain, swelling and impaired performance.⁽¹⁾ It is a common cause of posterior heel pain and is more prevalent in young and active populations, particularly athletes.⁽²⁾ The annual incidence of this disorder in top-level runners is 7%–9%.⁽³⁾ Although the exact pathogenesis of IAT remains unknown, increased mechanical strain, advancing age and decreased vascularity are often deemed as causative factors.⁽⁴⁻⁶⁾ Clinical presentations include localised pain over the posterior heel, swelling, burning and stiffness, which usually are aggravated by exertion.⁽⁷⁾ Radiology may show calcifications at the insertion of the Achilles tendon, with oedema and thickening of the insertional area of the tendon that may be seen on magnetic resonance imaging.^(8,9)

Nonoperative treatments, such as activity and footwear modification, eccentric training and extracorporeal shockwave therapy, are the primary management.⁽¹⁰⁻¹²⁾ Should nonoperative treatments fail, operative treatments may be considered. Surgery typically includes partial or total detachment of the Achilles tendon, excision of retrocalcaneal bursa, resection of prominent superior calcaneal tuberosity, debridement of the diseased tendon and reattachment of the tendon.^(6,12) A tendon transfer may be considered.^(4,13) The surgery can be done either by open or endoscopic means. Open surgery may be performed with a single longitudinal incision (placed either laterally or medially) or a central tendon-splitting incision. A single transverse incision, a J-shaped incision, and a combination of lateral and medial incisions have also been described. However, to the best of our knowledge, the most effective open operative approach has yet to be defined.

The objective of this study was to retrospectively compare the outcomes of operative treatment of IAT between the lateral approach (LA) and the central tendon-splitting approach (CA), using prospectively collected data.

METHODS

This study was approved by our institutional review board and informed consent was obtained. A retrospective review was done of patients who underwent operative treatment for IAT between August 2007 and May 2015 at Singapore General Hospital, Singapore. All patients were diagnosed with IAT based on preoperative clinical, imaging and intraoperative findings of the affected heel. The operative procedure included partial detachment of the Achilles tendon, debridement of the diseased tendon, excision of retrocalcaneal bursa, resection of prominent superior calcaneal tuberosity and reattachment of the tendon. Patients were excluded if they had a previous Achilles tendon rupture or other previous surgeries, had infection over the same heel, or had undergone operative correction with complete detachment of the Achilles tendon.

Patients were categorised into the LA group and CA group based on the two different open operative approaches. All patients' medical records were retrieved. Data extracted included gender, age, body mass index (BMI), length of conservative management, duration of surgery, length of hospitalisation and postoperative complications. Preoperative data that was prospectively collected included: visual analogue scale (VAS) scores to measure preoperative and postoperative subjective pain during activity (with 0 indicating no pain and 10 indicating extremely severe pain); American Orthopaedic Foot and Ankle Society (AOFAS) Ankle-Hindfoot Scale scores; and 36-item Short Form Health Survey (SF-36) scores to evaluate functional outcome. In the SF-36, eight parameters are scored on a scale of 0 to 100, with 100 being the maximal level of possible function. Scores of a subjective question involving postoperative satisfaction were also prospectively recorded. Patients were asked to rate the overall result of the surgery (1 = excellent, 2 = very good, 3 = good, 4 = fair, 5 = poor and 6 = terrible).

All operations were performed by the senior author (i.e. Rikhraj IS). Patients were given prophylactic antibiotics at the induction of anaesthesia and for 24 hours postoperatively, together with adequate analgesics. A thigh tourniquet was applied for all patients. For the lateral approach, as described by Lin et al,⁽¹⁴⁾ the patient was placed in the lateral position under anaesthesia. A lateral incision parallel to the lateral border of the Achilles tendon was marked out (Fig. 1) and made. The Achilles tendon was exposed and partially detached from the lateral aspect, and the diseased part of the tendon was debrided. For the central tendon-splitting approach, as described by McGarvey et al⁽¹⁵⁾ and Xia et al,⁽¹⁶⁾ the patient was placed in the prone position with a midline incision marked out (Fig. 2). The Achilles tendon was centrally split and partially detached from the calcaneum along the tendon insertion from the midline to the lateral half.

For both groups, the debridement of fibrous degeneration and calcification dictated the amount of tendon that remained. This was decided by the surgeon through repetitive palpation of the tendon. The retrocalcaneal bursa was excised to expose the superior aspect of the calcaneum, and Haglund's prominence was resected using an oscillating saw and rongeurs. Subsequently, the Achilles tendon was reattached to the posterior calcaneum using two suture anchors (HEALIX BRTM Anchor with Orthocord, Corkscrew FT II [Arthrex, Naples, FL, USA] or InsiteTM FT Suture Anchor System [Tornier, Bloomington, MN, USA]), with the foot in plantigrade position. The wound was thoroughly irrigated, followed by closure of subcutaneous tissue and the skin.

Sutures over the operative wound were removed between postoperative day (POD) 14 and POD 21 based on the condition of wound healing. All patients followed the same rehabilitation protocol. A non-weight-bearing plaster cast was applied with the foot in plantigrade position for 1–3 weeks after the operation, followed by a weight-bearing-as-tolerated controlled ankle motion walker boot until six weeks postoperatively. Footwear was

subsequently allowed. Following removal of the cast or the walker boot, rehabilitation was initiated, consisting of Achilles tendon stretching, ankle range of motion exercises and proprioception training. The rehabilitation was continued until functional goals were met. We prospectively recorded postoperative VAS, AOFAS Ankle-Hindfoot scale and SF-36 scores at the three-month postoperative visit and the last visit, as well as satisfaction rate at the last visit.

Statistical analysis was performed using IBM SPSS Statistics version 20.0 (IBM Corp, Armonk, NY, USA). Two-tailed paired *t*-test and one-way analysis of variation were used to assess the differences in preoperative and postoperative data. Mann-Whitney *U* test and chi-square test were applied to evaluate the differences between the two groups. The level of significance was set at $p < 0.05$.

RESULTS

The data of 84 patients who were diagnosed with insertional Achilles tendinopathy (i.e. 84 heels) and had undergone operative treatment were reviewed. 13 patients who had previous surgeries on the same heel were excluded. The remaining 71 patients were allocated into the LA group (39 heels) and CA group (32 heels). Patient demographics are summarised in Table I. There was no significant difference between the two groups in terms of age, gender, preoperative BMI, side of operated heel, operating duration and length of hospitalisation. The average follow-up duration was 16.1 ± 7.6 (range 10–45) months and 16.2 ± 4.9 (range 12–27) months in the LA group and CA group, respectively.

Table I. Demographic data of the patients.

Parameter	No. (%)		p-value
	LA group (n = 39)	CA group (n = 32)	
Age at surgery* (yr)	55 ± 12 (20.2–75.7)	58 ± 11 (25.9–78.9)	0.229
Gender			0.271
Female	28 (72)	19 (59)	
Male	11 (28)	13 (41)	
Body mass index [†]	28.1 ± 3.7	28.9 ± 3.9	0.367

Side of operated heel			
Right	23 (59)	19 (59)	0.973
Left	16 (41)	13 (41)	
Duration of surgery[†] (min)	44.3 ± 8.6	50.9 ± 20.3	0.446
Length of hospital stay[†] (day)	1.6 ± 1.3	1.7 ± 1.2	0.108
Follow-up duration* (mth)	16.1 ± 7.6 (10–45)	16.2 ± 4.9 (12–27)	0.283

Data is presented as *mean ± standard deviation (range), †mean ± standard deviation. LA: lateral approach; CA: central approach

In the LA group, mean VAS score improved significantly from 7.1 ± 1.7 preoperatively to 2.7 ± 2.5 at the three-month postoperative visit ($p < 0.001$) and 1.4 ± 2.1 at the last visit ($p < 0.001$). Similarly, in the CA group, the preoperative mean VAS score of 7.4 ± 1.7 also significantly improved to 4.4 ± 3.0 at three months postoperatively ($p < 0.001$) and to 0.7 ± 1.6 at the last visit ($p < 0.001$). When comparing VAS scores between the LA and CA groups, a significant difference was observed at the three-month postoperative visit ($p = 0.016$), while no significant difference was seen preoperatively ($p = 0.432$) and at the last visit ($p = 0.142$).

Average AOFAS Ankle-Hindfoot Scale scores in the LA group increased significantly from 43.1 ± 15.3 preoperatively to 68.5 ± 19.8 at the three-month postoperative visit ($p < 0.001$) and 88.7 ± 14.6 at the last visit ($p < 0.001$). In the CA group, scores increased significantly from 43.0 ± 17.6 preoperatively to 60.8 ± 21.2 at the three-month postoperative visit ($p < 0.001$) and 91.8 ± 9.8 at the last visit ($p < 0.001$). When comparing average AOFAS Ankle-Hindfoot Scale scores between the LA and CA groups, no significant difference was seen at the preoperative ($p = 0.945$), three-month postoperative ($p = 0.094$) or last visit ($p = 0.554$). SF-36 scores improved in most of the parameters, but there was no significant difference between the two groups postoperatively. These assessments are summarised in Table II.

Table II. Clinical outcomes of the patients.

Parameter	Mean ± standard deviation		p-value
	LA group	CA group	
VAS pain scale			

Preoperative	7.1 ± 1.7	7.4 ± 1.7	0.432
3-mth postoperative	2.7 ± 2.5	4.4 ± 3.0	0.016*
Last visit	1.4 ± 2.1	0.7 ± 1.6	0.142
p-value	< 0.001*	< 0.001*	
AOFAS Ankle-Hindfoot Scale			
Preoperative	43.1 ± 15.3	43.0 ± 17.6	0.945
3-mth postoperative	68.5 ± 19.8	60.8 ± 21.2	0.094
Last visit	88.7 ± 14.6	91.8 ± 9.8	0.554
p-value	< 0.001*	< 0.001*	
SF-36			
Physical function			
<i>Preoperative</i>	58.7 ± 23.0	53.0 ± 27.6	0.375
<i>Last visit</i>	78.0 ± 18.1	78.0 ± 15.0	0.678
<i>p-value</i>	< 0.001*	0.008*	
Role limitations (physical)			
<i>Preoperative</i>	24.4 ± 36.0	24.2 ± 38.4	0.840
<i>Last visit</i>	72.4 ± 39.7	76.6 ± 39.1	0.577
<i>p-value</i>	0.235	0.059	
Bodily pain			
<i>Preoperative</i>	34.2 ± 16.3	34.3 ± 21.4	0.753
<i>Last visit</i>	65.7 ± 26.6	67.1 ± 22.5	0.806
<i>p-value</i>	0.152	0.033*	
General health			
<i>Preoperative</i>	67.5 ± 27.7	76.2 ± 17.4	0.330
<i>Last visit</i>	73.3 ± 20.2	72.6 ± 20.3	0.949
<i>p-value</i>	0.364	0.015*	
Vitality			
<i>Preoperative</i>	66.4 ± 24.5	68.9 ± 21.2	0.735
<i>Last visit</i>	72.8 ± 20.9	73.3 ± 19.2	0.883
<i>p-value</i>	0.003*	0.003*	
Social function			
<i>Preoperative</i>	65.7 ± 36.6	59.4 ± 38.6	0.493
<i>Last visit</i>	88.1 ± 26.4	96.9 ± 13.8	0.084
<i>p-value</i>	< 0.001*	0.176	
Role limitations (emotional)			
<i>Preoperative</i>	84.6 ± 32.3	96.9 ± 17.7	0.034*
<i>Last visit</i>	96.6 ± 14.9	98.4 ± 8.8	0.654
<i>p-value</i>	0.121	0.861	
Mental health			
<i>Preoperative</i>	76.7 ± 18.1	79.3 ± 19.8	0.402
<i>Last visit</i>	84.7 ± 12.3	85.9 ± 13.1	0.443
<i>p-value</i>	0.044*	< 0.001*	

*Difference is statistically significant ($p < 0.05$). AOFAS: American Orthopaedic Foot and Ankle Society; CA: central approach; LA: lateral approach; SF-36: 36-item Short Form Health Survey; VAS: visual analogue scale

Postoperative complications for both study groups are summarised in Table III. Complications in the LA group included painful scarring (7.7%), wound sensation impairment (7.7%), delayed wound healing (2.6%), superficial wound infection (2.6%) and overgrown osteophytes at the heel requiring a secondary operation (2.6%), while those in the CA group included painful scarring (6.3%), wound sensation impairment (3.1%) and delayed wound healing (15.6%). There was no superficial wound infection in the CA group. No Achilles tendon rupture or deep wound infection was noted in both groups. Delayed wound healing, which occurs when a wound takes a longer-than-expected time to completely heal, was defined in this study as complete healing of a heel wound that takes longer than three weeks. We observed that the occurrence of delayed wound healing in the CA group was significantly higher as compared with the LA group ($p = 0.049$). For all other complications, no significant difference was detected. The overall complication rate was comparable, at 23.1% in the LA group and 25.0% in the CA group ($p = 0.850$). The overall satisfaction rate was 87.2% in the LA group and 87.5% in the CA group ($p = 0.968$).

Table III. Postoperative complications and satisfaction rate.

Parameter	No. (%)		p-value
	LA group (n = 39)	CA group (n = 32)	
Postoperative complication			
Painful scar	3 (7.7)	2 (6.3)	0.813
Wound sensation impairment	3 (7.7)	1 (3.1)	0.406
Delayed wound healing	1 (2.6)	5 (15.6)	0.049
Superficial wound infection	1 (2.6)	0 (0)	0.362
Overgrown osteophyte	1 (2.6)	0 (0)	0.362
Overall complication rate	9 (23.1)	8 (25.0)	0.850
Satisfaction rate	34 (87.2)	28 (87.5)	0.968

CA: central approach; LA: lateral approach

DISCUSSION

Although different open approaches have been described for operative management of IAT, there is a paucity of literature on the comparison between different approaches, and the optimal

approach has not been defined. The current study reviewed 71 heels that were diagnosed with IAT and treated surgically, comparing the outcomes of 39 heels that were operated on using the LA with that of 32 heels for which the CA was used. At the best of our knowledge, this is the study with the largest sample size that focuses on comparing the effectiveness and complications of the LA and CA for IAT.

The LA to the posterior heel was popularised by Palmer⁽¹⁷⁾ and is a common approach for IAT. It provides direct exposure of the posterolateral heel, particularly superolateral calcaneal exostosis. The major disadvantage of this approach is sural nerve injury, leading to numbness or paraesthesia of the skin.⁽¹⁸⁾ Yodlowski et al⁽¹⁹⁾ reported that as many as 34.1% of the patients had altered sensation of the skin. However, the incidence of impaired wound sensation in their study was 7.7%, with no statistically significant difference in wound sensation impairment between the two approaches. Our relatively low rate of nerve injury may be explained by the fact that soft tissue was meticulously handled to minimise nerve injury.

The CA for IAT was first introduced by McGarvey et al in 2002⁽¹⁵⁾ and has gained more popularity in the past few years. It provides excellent visualisation of pathologies over the posterior heel and reduces the incidence of sural nerve injury.^(20,21) This approach also allows surgeons to treat intratendinous lesions or simultaneously perform a tendon release.^(22,23) A cadaveric study⁽²⁴⁾ showed that the central incision lies between the peroneal angiosome and the posterior tibial angiosome; this may minimise vascular compromise from the heel surgery. However, one of the drawbacks of the CA is that it results in substantial tension over the wound after closure and, therefore, may cause delayed wound healing and wound dehiscence, as were commonly reported in previous studies.^(15,20,21,25,26) Similarly, we observed in our study that the CA led to significantly more cases of delayed wound healing when compared with the LA.

Both the LA and CA have been demonstrated to be effective. In the LA group, there was significant improvement in pain (i.e. mean postoperative scores), AOFAS Ankle-Hindfoot

Scale scores and SF-36 scores in our study, and the findings were similar to those reported in previous studies.^(9,14,19,27) Similarly, significant improvement in pain and function were noted in the CA group, and the results were in agreement with observations reported by previous studies.^(20,25,28,29) The two approaches were comparable in effectiveness in improving pain scores, AOFAS Ankle-Hindfoot Scales and SF-36 scores, and this is consistent with the results of Anderson et al.⁽³⁰⁾ In our study, the satisfaction rate of surgery using the CA was 87.5%, which falls within the range of 77%–100% reported in the literature.^(15,28,29,31) The satisfaction rate with the LA was 87.2% with no significant difference between the two approaches in terms of subjective evaluation.

Interestingly, we observed that the CA resulted in significant more short-term pain when compared with the LA. However, both approaches were comparable in long-term pain relief. Although this may not be clinically significant, one possible explanation is that short-term heel pain may be related to the position and condition of the surgical wound, as recovery from open retrocalcaneal decompression is usually prolonged.^(20,27) The surgical wound from the CA lies posteriorly against the heel counter of footwear and tends to be more irritated by the pressure, particularly when the wound is still oedematous during recovery. However, once the patient has completely recovered, the wound from the CA does not cause more pain when compared to that from the LA.

The results of the current study need to be interpreted in light of its limitations. First, the follow-up duration is relatively short. Previous studies suggested that patients who underwent surgical correction for Haglund's deformity took six months to two years to recover.^(27,32) A longer follow-up duration may be required to evaluate the long-term outcomes of different approaches. Second, the number of patients in this study is relatively small, and the sample size may be not large enough to detect any potential difference between the two surgical approaches. Last, this is a retrospective study from a single institution, which may result in

selection and observational biases. A prospective multicentre study with a longer follow-up duration and larger sample size is required to further evaluate the advantages and disadvantages of the LA or CA for operative management of IAT.

In conclusion, the LA and the CA with the operative techniques we described were effective treatments for IAT that resulted in similar functional improvement, long-term pain relief and overall enhancement of patients' health conditions. The two approaches were comparable in overall postoperative complication rate. With meticulous soft tissue handling, the LA provided better pain relief in the short term and reduced delayed wound healing as compared with the CA.

REFERENCES

1. Maffulli N, Khan KM, Puddu G. Overuse tendon conditions: time to change a confusing terminology. *Arthroscopy* 1998; 14:840-3.
2. Rio E, Mayes S, Cook J. Heel pain: a practical approach. *Aust Fam Physician* 2015; 44:96-101.
3. Lysholm J, Wiklander J. Injuries in runners. *Am J Sports Med* 1987; 15:168-71.
4. Irwin T. Current concepts review: insertional achilles tendinopathy. *Foot Ankle Int* 2010; 31:933-9.
5. Bah I, Kwak ST, Chimenti RL, et al. Mechanical changes in the Achilles tendon due to insertional Achilles tendinopathy. *J Mech Behav Biomed Mater* 2016; 53:320-8.
6. Li HY, Hua YH. Achilles tendinopathy: current concepts about the basic science and clinical treatments. *Biomed Res Int* 2016; 2016:6492597.
7. Maffulli N, Sharma P, Luscombe KL. Achilles tendinopathy: aetiology and management. *J R Soc Med* 2004; 97:472-6.

8. Sundararajan PP, Wilde TS. Radiographic, clinical, and magnetic resonance imaging analysis of insertional Achilles tendinopathy. *J Foot Ankle Surg* 2014; 53:147-51.
9. Oshri Y, Palmanovich E, Brin YS, et al. Chronic insertional Achilles tendinopathy: surgical outcomes. *Muscles Ligaments Tendons J* 2012; 2:91-5.
10. Scott A, Huisman E, Khan K. Conservative treatment of chronic Achilles tendinopathy. *CMAJ* 2011; 183:1159-65.
11. Tu P, Bytomski J. Diagnosis of heel pain. *Am Fam Physician* 2011; 84:909-16.
12. Wiegerinck JI, Kerkhoffs GM, van Sterkenburg MN, Sierevelt IN, van Dijk CN. Treatment for insertional Achilles tendinopathy: a systematic review. *Knee Surg Sports Traumatol Arthrosc* 2013; 21:1345-55.
13. Witt BL, Hyer CF. Achilles tendon reattachment after surgical treatment of insertional tendinosis using the suture bridge technique: a case series. *J Foot Ankle Surg* 2012; 51:487-93.
14. Lin HA, Chong HA, Yeo W. Calcaneoplasty and reattachment of the Achilles tendon for insertional tendinopathy. *J Orthop Surg (Hong Kong)* 2014; 22:56-9.
15. McGarvey WC, Palumbo RC, Baxter DE, Leibman BD. Insertional Achilles tendinosis: surgical treatment through a central tendon splitting approach. *Foot Ankle Int* 2002; 23:19-25.
16. Xia Z, Yew AKS, Zhang TK, et al. Surgical correction of Haglund's triad using a central tendon-splitting approach: a retrospective outcomes study. *J Foot Ankle Surg* 2017; 56:1132-8.
17. Palmer I. The mechanism and treatment of fractures of the calcaneus; open reduction with the use of cancellous grafts. *J Bone Joint Surg Am* 1948; 30A:2-8.

18. Freeman BJ, Duff S, Allen PE, Nicholson HD, Atkins RM. The extended lateral approach to the hindfoot. Anatomical basis and surgical implications. *J Bone Joint Surg Br* 1998; 80:139-42.
19. Yodlowski ML, Scheller AD Jr, Minos L. Surgical treatment of Achilles tendinitis by decompression of the retrocalcaneal bursa and the superior calcaneal tuberosity. *Am J Sports Med* 2002; 30:318-21.
20. Johnson KW, Zalavras C, Thordarson DB. Surgical management of insertional calcific achilles tendinosis with a central tendon splitting approach. *Foot Ankle Int* 2006; 27:245-50.
21. Highlander P, Greenhagen RM. Wound complications with posterior midline and posterior medial leg incisions: a systematic review. *Foot Ankle Spec* 2011; 4:361-9.
22. Miao XD, Jiang H, Wu YP, et al. Treatment of calcified insertional Achilles tendinopathy by the posterior midline approach. *J Foot Ankle Surg* 2016; 55:529-34.
23. Oliva F, Via AG, Maffulli N. Physiopathology of intratendinous calcific deposition. *BMC Med* 2012; 10:95.
24. Attinger CE, Evans KK, Bulan E, Blume P, Cooper P. Angiosomes of the foot and ankle and clinical implications for limb salvage: reconstruction, incisions, and revascularization. *Plast Reconstr Surg* 2006; 117(7 Suppl):261S-93S.
25. Gillis CT, Lin JS. Use of a central splitting approach and near complete detachment for insertional calcific Achilles tendinopathy repaired with an Achilles bridging suture. *J Foot Ankle Surg* 2016; 55:235-9.
26. McAlister JE, Hyer CF. Safety of Achilles detachment and reattachment using a standard midline approach to insertional enthesophytes. *J Foot Ankle Surg* 2015; 54:214-9.

27. Brunner J, Anderson J, O'Malley M, et al. Physician and patient based outcomes following surgical resection of Haglund's deformity. *Acta Orthop Belg* 2005; 71:718-23.
28. Ettinger S, Razzaq R, Waizy H, et al. Operative treatment of the insertional Achilles tendinopathy through a transtendinous approach. *Foot Ankle Int* 2016; 37:288-93.
29. Ahn JH, Ahn CY, Byun CH, Kim YC. Operative treatment of Haglund syndrome with central Achilles tendon-splitting approach. *J Foot Ankle Surg* 2015; 54:1053-6.
30. Anderson JA, Suero E, O'Loughlin PF, Kennedy JG. Surgery for retrocalcaneal bursitis: a tendon-splitting versus a lateral approach. *Clin Orthop Relat Res* 2008; 466:1678-82.
31. Nunley JA, Ruskin G, Horst F. Long-term clinical outcomes following the central incision technique for insertional Achilles tendinopathy. *Foot Ankle Int* 2011; 32:850-5.
32. Schneider W, Niehus W, Knahr K. Haglund's syndrome: disappointing results following surgery -- a clinical and radiographic analysis. *Foot Ankle Int* 2000; 21:26-30.

FIGURES



Fig. 1 Photograph shows the lateral approach, parallel to the anterior border of the Achilles tendon, marked out on the heel of one of the subjects.



Fig. 2 Photograph shows the central tendon-splitting approach marked out on the heel of one of the subjects.