

Injury patterns in elderly cyclists and motorcyclists presenting to a tertiary trauma centre in Singapore

Hui Shyuan Cheong¹, MBBS, Kum Ying Tham², FRCSEd, MBBS, Li Qi Chiu², MMed, MBBS

INTRODUCTION With Singapore's ageing population, there are increasing numbers of elderly cyclists and motorcyclists. Compared to younger riders, this cohort sustains more injuries and has poorer outcomes. This study aimed to describe and compare patient demographics, injury patterns and outcomes among elderly cyclists and motorcyclists at a Level 1 trauma centre.

METHODS Data of all cyclists, motorcyclists and pillion riders aged 65 years and above who presented to the emergency department after accidents from 1 January 2013 to 31 December 2017 was extracted from the hospital's trauma registry and reviewed.

RESULTS Cyclists and motorcyclists formed 42.0% and 58.0%, respectively, of 157 recruited patients. At the time of the accident, 40.8% of the patients were employed. The mean age of the patients was 71.6 ± 5.8 years. Extremities and pelvic girdle injuries (61.1%) were the most frequent, followed by chest injuries (48.4%), and head and neck injuries (40.1%). Among severe injuries (defined as Abbreviated Injury Scale score ≥ 3), chest injuries (39.5%) were the most common, followed by head and neck injuries (36.3%). The overall mortality rate was 9.6%, with cyclists at nearly three times the risk compared to motorcyclists. More cyclists than motorcyclists (18.2% vs. 11.0%) required intensive care. There were no significant differences in the length of hospital stay between cyclists and motorcyclists.

CONCLUSION Elderly riders have unique injury patterns and consume significant healthcare resources. Trauma systems need to acknowledge this changing injury epidemiology and equip trauma centres with the necessary resources targeted at elderly patients. Future work should focus on strategies to minimise extremity and chest injuries.

Keywords: geriatric trauma, injury patterns, motorcycle accidents

INTRODUCTION

Singapore is a city-state with a high population density of 7,804 persons/km² and an ageing population, with nearly 13% of the population aged above 65 years.⁽¹⁾ During 2013–2017, national statistics showed a total of 21,415 motorcycle fatalities.⁽²⁾ Of these, 12% were associated with motorcyclists aged above 60 years.⁽²⁾ Motorcycles and bicycles are important and affordable modes of transport whose increasing popularity is also related to the advent of bicycle-sharing systems and a demand for food delivery services. Data from the Singapore Police Force and other local studies has shown that motorcyclists and cyclists are among the most vulnerable road users.^(3–5) Regionally, motorcyclists and cyclists (along with pedestrians) make up half of all road traffic deaths in Southeast Asia.⁽⁶⁾ Studies have shown that compared to younger riders, elderly motorcyclists sustain more injuries^(7–9) as well as more severe injuries,^(9–14) and show different injury patterns.^(9,14) The increasing age of motorcyclists correlated with prolonged hospital stay,^(9,10,14,15) increased complications,^(9,10,15) increased mortality^(10,14,15) and higher care demands after discharge.⁽⁹⁾ Similarly, for cyclists, older age was associated with increased injury severity, mortality and length of intensive care unit stay.^(16–19)

Current literature comparing cyclists and motorcyclists has revealed differences in injury patterns and outcomes.^(20–22) However, to our knowledge, no study has been done to evaluate injury patterns and outcomes specifically in elderly cyclists and motorcyclists. This study aimed to describe and compare the patterns of injuries in elderly cyclists and motorcyclists presenting

to the emergency department (ED) at Tan Tock Seng Hospital (TTSH), Singapore.

METHODS

This was a retrospective review of all electronic medical records of cyclists, motorcyclists and pillion riders aged 65 years and above who presented to the ED at TTSH from 1 January 2013 to 31 December 2017 following accidents. TTSH is a 1,600-bed Level 1 trauma centre situated in a region of high population density and older housing estates. It has the busiest ED in Singapore, with over 170,000 attendances annually.

Data was obtained from the hospital's trauma registry. Data collected included patient demographics, mode of conveyance to the ED, triage Patient Acuity Category (PAC) scores, injury characteristics and patient outcomes. Injury characteristics included the epidemiology and severity of injuries according to Abbreviated Injury Scale (AIS) scores, Injury Severity Score (ISS), Revised Trauma Score (RTS) and probability of survival (Ps) scores. The study was approved by the National Healthcare Group Domain Specific Review Board.

Analysis was done using IBM SPSS Statistics version 25.0 (IBM Corp, Armonk, NY, USA). Means and standard deviations, and medians and interquartile ranges were calculated for normal and non-normal variables, respectively. Frequencies were tabulated for categorical variables. Student's *t*-test and chi-square test were used for analysis. A *p*-value < 0.05 was considered statistically significant.

¹Lee Kong Chian School of Medicine, Nanyang Technological University, ²Department of Emergency Medicine, Tan Tock Seng Hospital, Singapore

Correspondence: Dr Chiu Li Qi, Consultant, Department of Emergency Medicine, Tan Tock Seng Hospital, 11 Jalan Tan Tock Seng Singapore 308433. chiuliqi@gmail.com

RESULTS

A total of 157 elderly cyclists and motorcyclists presented to our centre from 1 January 2013 to 31 December 2017. Of these, 42.0% (n = 66) were cyclists. The mean age of the patients was 71.6 ± 5.8 years. Most riders (94.3%) were men. At the time of the accident, 40.8% of the patients were employed. A majority (81.5%) of patients arrived at the ED via emergency ambulance services and were triaged as PAC 2 (64.3%). One-third (34.4%) of the patients were triaged as PAC 1 and managed in the resuscitation area. The characteristics of cyclists and motorcyclists were similar, except that a higher proportion of elderly cyclists were retired or unemployed (66.7%) when compared to motorcyclists (46.2%). Table I shows the demographics of the study population.

Table II shows the injury patterns categorised according to AIS regions. Excluding AIS region 6 (external system), injuries involving extremities and the pelvic girdle (61.1%) were the most frequent, followed by chest injuries (48.4%), and head and neck injuries (40.1%). Among severe injuries (defined as AIS score ≥ 3), chest injuries were the most common (39.5%), followed by head and neck injuries (36.3%), and extremity and pelvic girdle

Table I. Demographics of elderly cyclists and motorcyclists.

Variable	No. (%)		
	All (n = 157)	Cyclists (n = 66)	Motorcyclists (n = 91)
Age* (yr)	71.6 ± 5.8	73.0 ± 6.0	70.5 ± 5.4
Male gender	148 (94.3)	61 (92.4)	87 (95.6)
Mode of arrival at ED			
Emergency ambulance service	128 (81.5)	52 (78.8)	76 (83.5)
Private ambulance	10 (6.4)	4 (6.1)	6 (6.6)
Own transport	19 (12.1)	10 (15.2)	9 (9.9)
Employment status			
Employed	64 (40.8)	19 (28.8)	45 (49.5)
Retired/unemployed	86 (54.8)	44 (66.7)	42 (46.2)
Unknown	7 (4.5)	3 (4.5)	4 (4.4)
Triage PAC score			
PAC 1	54 (34.4)	20 (30.3)	34 (37.4)
PAC 2	101 (64.3)	45 (68.2)	56 (61.5)
PAC 3	2 (1.3)	1 (1.5)	1 (1.1)

*Data presented as mean ± standard deviation. ED: emergency department; PAC: Patient Acuity Category

Table II. Injury patterns of elderly cyclists and motorcyclists according to AIS regions.

Variable	No. (%) / mean ± standard deviation			p-value
	All (n = 157)	Cyclists (n = 66)	Motorcyclists (n = 91)	
Region of injury				
Head and neck	63 (40.1)	30 (45.5)	33 (36.3)	0.246
Face	25 (15.9)	10 (15.2)	15 (16.5)	0.822
Chest	76 (48.4)	26 (39.4)	50 (54.9)	0.045
Abdominal and pelvic contents	15 (9.6)	6 (9.1)	9 (9.9)	0.866
External	113 (72.0)	46 (69.7)	67 (73.6)	0.588
Extremities and pelvic girdle	96 (61.1)	34 (51.5)	62 (68.1)	0.035
<i>Upper extremity</i>	43 (27.4)	10 (15.2)	33 (36.3)	0.003
<i>Lower extremity</i>	62 (39.5)	27 (40.9)	35 (38.5)	0.757
Fracture site				
Scapula	11 (7.0)	2 (3.0)	9 (9.9)	NS
Clavicle	16 (10.2)	4 (6.1)	12 (13.2)	NS
Humerus	7 (4.5)	2 (3.0)	5 (5.5)	NS
Forearm	6 (3.8)	1 (1.5)	5 (5.5)	NS
Hand	7 (4.5)	1 (1.5)	6 (6.6)	NS
Pelvic ring	6 (3.8)	3 (4.5)	3 (3.3)	NS
Acetabulum	6 (3.8)	2 (3.0)	4 (4.4)	NS
Femur	38 (24.2)	22 (33.3)	16 (17.6)	0.023
Patella	3 (1.9)	1 (1.5)	2 (2.2)	NS
Tibia/fibula	16 (10.2)	2 (3.0)	14 (15.4)	0.012
Foot	3 (1.9)	0 (0)	3 (3.3)	NS
Injury severity				
ISS	16.1 ± 10.3	16.6 ± 10.6	15.7 ± 10.0	0.587
RTS	7.5 ± 1.0	7.4 ± 0.9	7.6 ± 1.1	0.505
Ps	87.7 ± 20.2	85.2 ± 21.7	89.5 ± 18.9	0.185
ISS ≥ 16	51 (32.5)	23 (34.8)	28 (30.8)	0.590

AIS: Abbreviated Injury Scale; ISS: Injury Severity Score; NS: not significant; Ps: probability of survival; RTS: Revised Trauma Score

injuries (34.4%). The occurrence of severe chest injuries ($p = 0.045$), and extremity and pelvic girdle injuries ($p = 0.035$) among motorcyclists were significantly higher than among cyclists. Among extremity injuries, femoral fractures (24.2%) were the most common. A significantly higher proportion of motorcyclists sustained upper extremity injuries when compared to cyclists ($p = 0.003$). Cyclists sustained significantly more femoral fractures than motorcyclists ($p = 0.023$). The reverse was observed for tibia/fibular fractures ($p = 0.012$). There were no significant differences in the mean ISS, RTS and Ps scores between cyclists and in the motorcyclists, and in the proportion of patients with major trauma (defined as $ISS \geq 16$).

Table III shows the outcomes of our patients. One patient succumbed to injuries within the ED. All other patients required hospital admission, with a median length of stay of 6.0 (range 2.0–10.8) days. The overall mortality rate was 9.6%, with cyclists having nearly three times the mortality risk of motorcyclists. All of the 15 patients who died were men and had an average age of 74 years, and one-third were motorcyclists. The average length of stay before death was 3.6 days, with 11 patients dying within the first day of admission, three patients dying within the first week and one patient dying after a prolonged stay of one month. Head injuries ($n = 9$) were the overwhelming cause of death, with other reasons being head and chest injuries ($n = 2$), multiple injuries ($n = 2$) and blunt abdominopelvic injuries ($n = 1$). The patient who died at the one-month mark succumbed to pneumonia and renal failure subsequent to head injury. Among the survivors, one-third of both cyclists and motorcyclists required admission to the intensive care unit or high-dependency unit, while most of the others were admitted to the general ward. More cyclists than motorcyclists (18.2% vs. 11.0%) required intensive care. There were no significant differences in the length of hospital stay and disposition status among these patients. A majority (55.4%) of patients were discharged home and 28.0% of patients were discharged to a rehabilitation facility.

DISCUSSION

To our knowledge, this is the first study that compares injury patterns among elderly cyclists and motorcyclists. According to local statistics, elderly motorcyclists made up approximately 12% of all traffic casualties during the study period.⁽²⁾ Our results showed that 40.8% of patients were employed at the time of their accident. This suggests that despite the prevailing retirement age of 62 years, a large proportion of the elderly remain actively employed and contribute to the economy. Accidents in this cohort thus represent potential societal economic losses.

Care for this group consumes more healthcare resources than that for younger patients, as evidenced by one-third of our patients requiring management in the resuscitation area with subsequent admission to the intensive care unit and high-dependency unit. More than a quarter of our patients were discharged to a rehabilitation facility. Unfortunately, longitudinal data on functional outcomes was incomplete, and the impact on long-term outcomes among our patients could not be determined. Nevertheless, our findings

Table III. Outcomes of elderly cyclists and motorcyclists.

Patient outcome	No. (%)		
	All (n = 157)	Cyclists (n = 66)	Motorcyclists (n = 91)
Disposition from ED			
Intensive care unit	22 (14.0)	12 (18.2)	10 (11.0)
High dependence unit	32 (20.4)	12 (18.2)	20 (22.0)
Surgical acute care unit	1 (0.6)	0 (0)	1 (1.1)
General ward	101 (64.3)	42 (63.6)	59 (64.8)
Death within ED	1 (0.6)	0 (0)	1 (1.1)
Length of hospital stay* (day)	6 (2.0–10.8)	6.5 (2.0–10.3)	5 (2.0–11.0)
Length of ICU stay* (day)	2 (1.0–4.0)	1 (1.0–3.0)	2 (1.0–7.0)
Discharge location			
Home	87 (55.4)	27 (40.9)	60 (65.9)
Rehabilitation facility	44 (28.0)	23 (34.8)	21 (23.1)
Nursing home	6 (3.8)	3 (4.5)	3 (3.3)
Discharged against medical advice	5 (3.2)	3 (4.5)	2 (2.2)
In-hospital mortality	15 (9.6)	10 (15.2)	5 (5.5)

*Data presented as median (interquartile range). ICU: intensive care unit; ED: emergency department

emphasised the impact older riders have on healthcare resource utilisation and were similar to the findings of other studies.^(9,18)

Results revealed that among elderly cyclists and motorcyclists, the extremities and pelvic girdle region were the most vulnerable to injury. This was consistent with previous studies involving cyclists^(18,19,23-27) and motorcyclists.^(11,27,28) However, the patterns of extremity injuries differed between cyclists and motorcyclists in our study. Motorcyclists sustained significantly higher rates of upper extremity injuries and tibia/fibular fractures, while cyclists sustained significantly more femoral fractures. Apart from Liu et al,⁽²⁰⁾ who found that cyclists had a higher odds ratio for femoral fractures when compared to motorcyclists, few studies have been done to investigate these differences. Further studies focusing on injury mechanisms are needed. Unlike head injuries, which may be reduced by helmet use, little is known about prevention of extremity injuries. Studies have shown that extremity injuries have high healthcare and productivity costs,^(29,30) and are negatively associated with the quality of well-being outcome.^(31,32) Resources should be allocated to investigate ways to minimise extremity injuries in order to reduce associated morbidity and economic costs.

The occurrence of severe chest injuries was significantly higher in elderly motorcyclists than in cyclists. Thoracic injuries are clinically significant because they are more likely to be associated with fatal outcomes,^(7,33) and this risk increases with age.⁽¹²⁾ As the area of thoracic exposure is comparable, a possible explanation could be that motorcyclists sustain greater impact during accidents due to higher speeds. This has a significant bearing on injury

prevention for elderly motorcyclists. Possible strategies to explore include stricter laws regarding speed limits and protective gear.

It is well established that helmets reduce the frequency and severity of head injuries in riders. Current traffic laws in Singapore do not mandate the use of helmets for cyclists. Interestingly, our study did not show any significant differences in head injuries between cyclists and motorcyclists. However, as this was not an *a priori* analysis, further studies specifically designed to evaluate the utility of helmets for head injury prevention in our population are necessary.

There were some limitations to this study. It was a retrospective, single-centre study with a small sample size. It omitted data from riders who did not present to the ED or died at the scene of the accident. Although it can be assumed that persons who did not present to the ED probably had more minor injuries, injury patterns among the most severely injured patients in our study were not captured. Longitudinal data regarding functional capacity after discharge was also unavailable for comparison due to these patients being lost to follow-up.

In conclusion, elderly motorcyclists and cyclists have unique injury patterns and consume significant healthcare resources. As our population ages and two-wheeled vehicles continue to be an indispensable mode of transport, this burden of injury will only increase. Trauma systems need to acknowledge this changing injury epidemiology and equip trauma centres with the necessary resources targeted at elderly riders. Future work should focus on strategies to minimise extremity and chest injuries. A prospective multicentre study using the nation's trauma database would be useful to assess the national burden of geriatric trauma among cyclists and motorcyclists. In addition, such a study should address possible differences in epidemiology and outcomes among younger riders when compared to older ones.

REFERENCES

- Department of Statistics, Singapore. Population and Population Structure. Available at: <https://www.singstat.gov.sg/find-data/search-by-theme/population/population-and-population-structure/latest-data>. Accessed January 4, 2019.
- Singapore Police Force. Traffic Annual 2017. In: Publication [online]. Available at: <https://www.police.gov.sg/news-and-publications/publications?category=Annual%20Reports>. Accessed December 20, 2018.
- Wong ZH, Chong CK, Tai BC, Lau G. A review of fatal road traffic accidents in Singapore from 2000 to 2004. *Ann Acad Med Singapore* 2009; 38:594-6.
- Wong E, Leong MK, Anantharaman V, et al. Road traffic accident mortality in Singapore. *J Emerg Med* 2002; 22:139-46.
- World Health Organization. Road Safety in the South-East Asian Region 2015. Available at: http://www.who.int/violence_injury_prevention/road_safety_status/2015/Road_Safety_SEAR_3_for_web.pdf. Accessed December 20, 2018.
- Department of Statistics, Singapore. Population Overview. Available at: <https://www.singstat.gov.sg/find-data/search-by-theme/population/population-and-population-structure/visualising-data/population-trends>. Accessed May 6, 2018.
- Dischinger PC, Ryb GE, Ho SM, Braver ER. Injury patterns and severity among hospitalised motorcyclists: a comparison of younger and older riders. *Annu Proc Assoc Adv Automot Med* 2006; 50:237-49.
- Ashie A, Wilhelm A, Carney D, DiPasquale T, Bush C. Comparing fracture patterns of younger versus older riders involved in nonfatal motorcycle accidents. *Traffic Inj Prev* 2018; 19:761-5.
- Muratore S, Hawes L, Farhat J, et al. Riding into the golden years: injury patterns and outcomes of advanced-age motorcycle trauma. *Am J Surg* 2016; 212:670-6.
- Brown JB, Bankey PE, Gorczyca JT, et al. The aging road warrior: national trend toward older riders impacts outcome after motorcycle injury. *Am Surg* 2010; 76:279-86.
- Jackson TL, Mello MJ. Injury patterns and severity among motorcyclists treated in US emergency departments, 2001-2008: a comparison of younger and older riders. *Inj Prev* 2013; 19:297-302.
- Talving P, Teixeira PG, Barmparas G, et al. Motorcycle-related injuries: effect of age on type and severity of injuries and mortality. *J Trauma* 2010; 68:441-6.
- Stutts J, Foss R, Svoboda C. Characteristics of older motorcyclist crashes. *Annu Proc Assoc Adv Automot Med* 2004; 48:197-211.
- Hsieh CH, Liu HT, Hsu SY, Hsieh HY, Chen YC. Motorcycle-related hospitalizations of the elderly. *Biomed J* 2017; 40:121-8.
- Warlick B, Nuismer A, Israel H, Cannada LK. The aging motorcyclist: a comparative epidemiologic study on pattern and severity of injury. *J Trauma Acute Care Surg* 2012; 73:175-8.
- Chong S, Poulos R, Olivier J, Watson WL, Grzebieta R. Relative injury severity among vulnerable non-motorised road users: comparative analysis of injury arising from bicycle-motor vehicle and bicycle-pedestrian collisions. *Accid Anal Prev* 2010; 42:290-6.
- Cripton PA, Shen H, Brubacher JR, et al. Severity of urban cycling injuries and the relationship with personal, trip, route and crash characteristics: analyses using four severity metrics. *BMJ Open* 2015; 5:e006654.
- Lustenberger T, Inaba K, Talving P, et al. Bicyclists injured by automobiles: relationship of age to injury type and severity--a national trauma databank analysis. *J Trauma* 2010; 69:1120-5.
- Rivara FP, Thompson DC, Thompson RS. Epidemiology of bicycle injuries and risk factors for serious injury. *Inj Prev* 1997; 3:110-4.
- Liu HT, Rau CS, Liang CC, et al. Bicycle-related hospitalizations at a Taiwanese level I Trauma Center. *BMC Public Health* 2015; 15:722.
- Pai CW, Chen YC, Lin HY, Chen PL. A population-based case-control study of hospitalisation due to head injuries among bicyclists and motorcyclists in Taiwan. *BMJ Open* 2017; 7:e018574.
- Pai CW, Lin HY, Tsai SH, Chen PL. Comparison of traffic-injury related hospitalisation between bicyclists and motorcyclists in Taiwan. *PLoS One* 2018; 13:e0191221.
- Davidson JA. Epidemiology and outcome of bicycle injuries presenting to an emergency department in the United Kingdom. *Eur J Emerg Med* 2005; 12:24-9.
- Dinh MM, Kastelein C, Hopkins R, et al. Mechanisms, injuries and helmet use in cyclists presenting to an inner city emergency department. *Emerg Med Australas* 2015; 27:323-7.
- Neumann MV, Eley R, Vallmuur K, Schuetz M. Current profile of cycling injuries: a retrospective analysis of a trauma centre level 1 in Queensland. *Emerg Med Australas* 2016; 28:90-5.
- Gopinath B, Jagnoor J, Craig A, et al. Describing and comparing the characteristics of injured bicyclists and other injured road users: a prospective cohort study. *BMC Public Health* 2016; 16:324.
- Chichom-Mefire A, Atashili J, Tsiagadigui JG, Fon-Awah C, Ngowe-Ngowe M. A prospective pilot cohort analysis of crash characteristics and pattern of injuries in riders and pillion passengers involved in motorcycle crashes in an urban area in Cameroon: lessons for prevention. *BMC Public Health* 2015; 15:915.
- Tham KY, Seow E, Lau G. Pattern of injuries in helmeted motorcyclists in Singapore. *Emerg Med J* 2004; 21:478-82.
- Meerding WJ, Looman CW, Essink-Bot ML, et al. Distribution and determinants of health and work status in a comprehensive population of injury patients. *J Trauma* 2004; 56:150-61.
- Polinder S, Haagsma J, Panneman M, et al. The economic burden of injury: health care and productivity costs of injuries in the Netherlands. *Accid Anal Prev* 2016; 93:92-100.
- Holbrook TL, Anderson JP, Sieber WJ, Browner D, Hoyt DB. Outcome after major trauma: discharge and 6-month follow-up results from the Trauma Recovery Project. *J Trauma* 1998; 45:315-24.
- Holbrook TL, Anderson JP, Sieber WJ, Browner D, Hoyt DB. Outcome after major trauma: 12-month and 18-month follow-up results from the Trauma Recovery Project. *J Trauma* 1999; 46:765-73.
- Doyle D, Muir M, Chinn B. Motorcycle accidents in Strathclyde Region, Scotland during 1992: a study of the injuries sustained. *Health Bull (Edinb)* 1995; 53:386-94.



About the First Author

Dr Cheong Hui Shyuan is a medical officer who is currently still exploring her interests. One year into working life, she still enjoys her work and interacting with patients. In her free time, she enjoys spending precious time with family and friends, and a good yoga or spin session. She is fervently hoping for the day travel restrictions are lifted and she can go hiking again.