

# Safety, tolerability and efficacy of LEGA-Kid® mechanical percussion device versus conventional chest physiotherapy in children: a randomised, single-blind controlled study

Yuen Ling Hue<sup>1</sup>, MBBS, Lucy Chai See Lum<sup>2</sup>, MBBS, MRCP, Siti Hawa Ahmad<sup>3</sup>, Dip PT, Soon Sin Tan<sup>1</sup>, MBBS, Shin Yee Wong<sup>2</sup>, BSc, PhD, Anna Marie Nathan<sup>2</sup>, MBBCh, MRCPCH, Kah Peng Eg<sup>2</sup>, MBBS, MPaeds, Melissa de Bruyne Ming May Choon<sup>1</sup>, MB BCh BAO

**INTRODUCTION** Chest physiotherapy (CPT) may benefit children aged below five years who suffer from lower respiratory tract infection (LRTI). However, its effects depend on the technique used. This study aimed to determine whether mechanical CPT using the LEGA-Kid® mechanical percussion device is superior to manual CPT in children with LRTI.

**METHODS** Children aged five months to five years who were admitted and referred for CPT from January to April 2017 were randomised to either manual CPT or mechanical CPT with LEGA-Kid. Outcomes measured before intervention and two hours after intervention were respiratory rate (RR), oxygen saturation and modified Respiratory Distress Assessment Instrument (mRDAI) score.

**RESULTS** All 30 enrolled patients showed significant reduction in post-intervention RR and mRDAI scores. There was an 8% reduction in RR for the manual CPT group ( $p = 0.002$ ) and a 16.5% reduction in the mechanical CPT group ( $p = 0.0001$ ), with a significantly greater reduction in the latter ( $p = 0.024$ ). mRDAI scores decreased by 2.96 in the manual group ( $p = 0.0001$ ) and 3.62 in the mechanical group ( $p = 0.002$ ), with no significant difference between the groups. There was no significant improvement in oxygen saturation, and no adverse events were observed after CPT.

**CONCLUSION** Children receiving both manual and mechanical CPT showed improvements in respiratory distress symptoms, with no adverse effects. A combined strategy of nebulised hypertonic saline followed by CPT for LRTI removes airway secretions and results in improvements in moderately severe respiratory distress. The LEGA-Kid mechanical CPT method is superior to manual CPT in reducing the RR.

*Keywords: children, lower respiratory tract infections, manual chest physiotherapy, mechanical percussor, nebulised hypertonic saline*

## INTRODUCTION

Lower respiratory tract infections (LRTIs) are common in children aged below five years and are a leading cause of hospital admissions. LRTIs are associated with increased morbidity and mortality.<sup>(1)</sup> Most infections are viral in origin and associated with copious thick secretions. As a result, the airway becomes obstructed, resulting in severe respiratory distress and ventilation-perfusion mismatch.<sup>(2)</sup> The presence of airway oedema also increases airway resistance, thereby worsening hypoxia. The consequences of mechanical obstruction include reduced tidal volume, increased work of breathing and retained secretions, which further exacerbate respiratory distress.

Chest physiotherapy (CPT) assists in mucociliary clearance, which reduces airway resistance and improves ventilation.<sup>(3)</sup> Newer CPT methods, such as forced exhalation with an open glottis<sup>(4)</sup> and slow-flow techniques,<sup>(5,6)</sup> have shown short-term benefits to respiratory symptoms from bronchial obstruction, without impact on duration of hospitalisation. In patients who are hospitalised with acute bronchiolitis, conventional CPT and expiratory acceleration flow techniques have shown short-term

improvements in respiratory distress scores and positive parental satisfaction.<sup>(7)</sup> Conventional CPT and prolonged slow expiration techniques were effective in reducing clinical scores 48 hours after intervention when compared to only upper airway suctioning in patients with respiratory syncytial virus bronchiolitis of moderate severity.<sup>(8)</sup> However, its use has been limited because CPT requires an experienced therapist to perform.

A Cochrane analysis of 12 randomised controlled trials in 2016 compared physiotherapy with no intervention in patients with mild, moderate and severe acute bronchiolitis.<sup>(9)</sup> Physiotherapy techniques included conventional percussion with vibration plus postural drainage and passive flow-orientated expiratory techniques. Both techniques failed to improve severity status and time to recovery. However, one trial observed a small improvement in the Wang clinical score immediately after the intervention in patients with moderately severe acute bronchiolitis.<sup>(8)</sup> The authors of the review concluded that these physiotherapy techniques could not be used as standard clinical practice for hospitalised patients with severe bronchiolitis, and instead recommended exploring the combination of CPT with

<sup>1</sup>University Malaya Paediatric and Child Health Research Group, <sup>2</sup>Department of Paediatrics, University of Malaya, <sup>3</sup>Physiotherapy Division, Department of Rehabilitation Medicine, University of Malaya Medical Centre, Kuala Lumpur, Malaysia

**Correspondence:** Dr Lucy Lum Chai See, Professor, Department of Paediatrics, University of Malaya, 50603 Kuala Lumpur, Malaysia. lumcs@ummc.edu.my

salbutamol or hypertonic saline in patients with mild to moderate acute bronchiolitis.<sup>(9)</sup>

Effective CPT techniques require skilled personnel and may be difficult to perform in young children. LEGA, a mechanical device designed by Formedic Technology Sdn Bhd, Malaysia, has been used as an aid for CPT. This machine is a handheld, battery-operated percussor that provides clapping to the chest wall to mimic manual hand percussion. A randomised controlled cross-over study evaluating the efficacy and safety of LEGA compared to conventional CPT in adults with chronic obstructive pulmonary disease showed the device to be effective in generating cough and expectoration of sputum. The device was well tolerated and patients were able to use it independent of a chest physiotherapist.<sup>(10)</sup>

A similar but smaller device (LEGA-Kid®) has been designed by Formedic Technologies for use in children weighing 5–15 kg. It produces rotational movements with a predetermined frequency that creates percussion and vibration forces when in contact with the chest wall. The generated force travels through the chest wall and helps to loosen secretions. We hypothesised that LEGA-Kid would be as effective as manual CPT for mucus clearance, thereby reducing respiratory distress without causing significant patient distress. This study sought to compare the safety, tolerability and efficacy of LEGA-Kid with that of manual CPT in children with LRTI.

## METHODS

This single-blind, randomised controlled study was conducted in the paediatric wards of University Malaya Medical Centre (UMMC). Eligible children were randomly allocated to manual (control group) or mechanical CPT (study group) after informed consent was obtained. Approval from the UMMC Medical Ethics Committee was obtained prior to the start of the study (ID no. 20166-2502). The study was registered with the National Medical Research Registry (NMRR) of Malaysia (NMRR-16-2803-32022).

Children aged five months to five years, weighing between 5 kg and 15 kg, who were admitted with symptoms of LRTI and referred for CPT by the managing doctors were included into the study. Children were excluded if they had severe respiratory distress or heart failure, bone disease, thrombocytopenia or a modified Respiratory Distress Assessment Instrument (mRDAI) score  $\leq 6$  (Table I).

Patients were randomised to either manual or mechanical CPT using an online software, Research Randomizer 4.0,<sup>(11)</sup> in a ratio of 1:1, using permuted blocks of 4. The number '1' was assigned to the first eligible participant, the number '2' to the second, and so on till number '30'. This was a single-blinded study, where the intervention arm was known to the participants and the physiotherapist, but not to the physician performing the pre- and post-intervention evaluations.

Nebulised 3% hypertonic saline (4 mL over 15 minutes) was administered to all patients before the physiotherapy session. CPT, together with postural drainage, was performed by an experienced paediatric chest physiotherapist. The following steps were incorporated: (a) an assessment of the patient's condition,

vital signs and auscultation to identify the involved lung segment; (b) placement of the patient according to the postural drainage positions; (c) chest percussion and vibration manually or with LEGA-Kid device on the chest area; (d) removal or expectoration of mucus by gentle oral/nasal suction and provoked cough; and (e) technique with localised breathing to enhance expansion of the specific lung segment.

Mechanical CPT involved application of the percussor firmly and at right angles to the chest wall to make the voice quiver. The selected frequency was indicated by the numbers 1 to 10; the lower numbers signified percussions and higher numbers signified vibrations. Each physiotherapy session lasted for 15–20 minutes, during which the patients were shown their favourite videos and toys. Post CPT, positioning was performed to improve chest expansion. Oxygen therapy was continued as at  $T_0$ , pre-intervention. Bronchodilator therapy was avoided during the intervention period, which was about three hours.

All patients underwent a baseline evaluation at enrolment, and pre-intervention ( $T_0$ ) and two-hour post-intervention ( $T_{120}$ ) evaluations by a trained physician who was blinded to the method of CPT. The primary outcome measures evaluated were temperature, heart rate, RR, pulse oximetry with a Nellcor oximeter after being in room air for five minutes and modified Respiratory Distress Assessment Instrument (mRDAI). The mRDAI is a modification of the RDAI, which rates wheezing and respiratory distress on a scale of 0 to 17. The addition of assessment of crackles gives a total score of 23 (Table I). Both RDAI and mRDAI have shown good inter-observer reliability.<sup>(12)</sup> The Wang severity score,<sup>(13)</sup> which assesses wheezing, RR, retractions and general condition, was evaluated daily. Each clinical sign was scored from 0 to 3 (possible total range of scores: 0–12), with a higher score indicating more severe disease. The Wang clinical severity score has been reported to have good inter-rater reliability, with an intraclass correlation coefficient of 0.99.<sup>(14)</sup>

During the intervention, a research assistant monitored the heart rate and oxygen saturation and also looked out for any adverse events such as bradycardia, desaturation and vomiting. The secondary outcome measures were length of hospital stay and FLACC (Face, Legs, Activity, Cry, Consolability) scale to assess the level of distress<sup>(15)</sup> at  $T_0$ , during intervention ( $T_{CPT}$ ) and at  $T_{120}$ . The FLACC scale evaluates pain and discomfort in the paediatric age group. It has a score of 0 to 10, where 0 indicates relaxed and comfortable, 1–3 indicates mild distress, 4–6 indicates moderate pain and 7–10 indicates severe distress.

CPT was carried out daily until the mRDAI score was  $\leq 6$  or until discharge from hospital. Typical care involving the type of respiratory support and oxygen supplementation was determined by the ward clinicians, who were blinded to the method of CPT. Oxygen therapy was delivered via nasal cannula or high flow nasal cannula. Target oxygen saturation was 95%, while the choice of oxygen delivery was tailored to the patient's RR and the presence of retractions.

Data was entered using IBM SPSS Statistics version 23.0 (IBM Corp, Armonk, NY, USA). To reach the statistical effect of 80%

**Table I. Modified Respiratory Distress Assessment Instrument scale: wheezing, crackles and retractions.**

Variable	Point				Maximum point	
	0	1	2	3		
<b>Wheezing</b>						
Expiration	None	End	½	¾	All	4
Inspiration	None	Part	All			2
Location	None	Segmental: ≤ 2 of 4 lung fields	Diffuse: ≥ 3 of 4 lung fields			2
<b>Crackles</b>						
Expiration	None	Part	All			2
Inspiration	None	Part	All			2
Location	None	Segmental: ≤ 2 of 4 lung fields	Diffuse: ≥ 3 of 4 lung fields			2
<b>Retraction</b>						
Supraclavicular	None	Mild	Moderate	Marked		3
Intercostal	None	Mild	Moderate	Marked		3
Subcostal	None	Mild	Moderate	Marked		3

Within each variable, the subscores are summed to give a total score. The maximum total points are 8 for wheezing, 6 for crackles and 9 for retraction.


power and 5% significance, a sample size of 30 was required – 15 in the manual CPT arm and 15 in the mechanical arm (LEGA-Kid). Non-parametric data was analysed using the Mann-Whitney *U* test, and parametric data was analysed using paired and Student's *t*-tests. Paired *t*-test was used to compare pre- and post-intervention observations in each group, while Student's *t*-test was used to compare the difference between the groups. Chi-square test was used for nominal data. One-way analysis of variance for repeated measures was used to compare the daily evolution of the Wang scores. A *p*-value < 0.05 was considered statistically significant.

## RESULTS

A total of 30 patients were enrolled and randomised to receive either manual or mechanical CPT between 9 January 2017 and 30 April 2017. The patients completed 45 sessions in the manual group and 42 sessions in the mechanical group for a total of 87 physiotherapy sessions. The demographic data and baseline clinical characteristics of the patients are shown in Table II. The median mRDAI score and Wang clinical severity score were similar (*p* = 0.683) in both arms; both groups had similar severity of illness. A median mRDAI score of 11 at enrolment indicated moderately severe respiratory distress. A total of 12 patients (six in each arm) required oxygen therapy or respiratory support during the hospitalisation period.

Patients in both the manual and mechanical groups showed significant reductions in RR and mRDAI score at  $T_{120}$  compared to  $T_0$  (Table III). The mean reduction in RR was 3.6 breaths per minute (8% reduction, *p* = 0.002) in the manual group and 7.0 breaths per minute (16.5% reduction, *p* = 0.0001) in the mechanical group. The reduction in RR was significantly greater in the mechanical group (*p* = 0.024).

Similarly, there was a significant decrease in mRDAI scores between  $T_0$  and  $T_{120}$ : 2.96 points (27.2% reduction, *p* = 0.0001) in the manual group and 3.62 points (34.5% reduction, *p* = 0.0001) in the mechanical group. Similar improvements were observed in all three components of the mRDAI score at  $T_{120}$  compared to  $T_0$ .



**Fig. 1** Graph shows daily pre-intervention ( $T_0$ ) Wang clinical severity scores.

However, the reduction in mRDAI scores in the mechanical and manual groups was not significant (*p* = 0.144). Children in the mechanical group showed significant improvement in oxygen saturation at  $T_{120}$  (*p* = 0.013) compared to the manual group (*p* = 0.123). However, the improvement in oxygen saturation in the mechanical and manual groups was not significant (*p* = 0.0509).

Table IV shows the FLACC scores before ( $T_0$ ), during ( $T_{CPT}$ ) and after ( $T_{120}$ ) intervention. For both groups of patients, the mean FLACC score did not differ considerably at  $T_{CPT}$  (manual 3.80 vs. mechanical 3.71, *p* = 0.86) compared to that at  $T_0$  (3.6 vs. 3.3, *p* = 0.64). The mean change in FLACC score between  $T_0$  and  $T_{CPT}$  was 0.24 (*p* = 0.175) in the manual group and 0.43 (*p* = 0.06) in the mechanical group. The difference between these two groups was not significant (*p* = 0.503). However, a significant reduction in the  $T_{120}$  FLACC score was observed in both groups of patients (manual 2.96, *p* = 0.0001 vs. mechanical 2.71, *p* = 0.0001), although the difference between the groups was not significant.

The daily mean Wang scores decreased significantly day by day from enrolment till Day 4 in both groups, from 8.4 to 4 (*p* < 0.001) in the manual group and from 8.6 to 4.5 (*p* < 0.001) in the mechanical group (Fig. 1). The number of patients on oxygen supplementation or respiratory support decreased from 12 (six in

Table II. Demographic data and parameters at enrolment.

Variable	Median (IQR)/No. (%)		p-value
	Manual CPT (n = 15)	Mechanical CPT (n = 15)	
Age (mth)	11 (7–19)	10 (8–13)	0.884
Male gender	10 (66.7)	13 (86.7)	
Weight (kg)	8.6 (7.4–9.2)	8.1 (7.3–9.2)	0.693
<b>Diagnosis</b>			
Acute bronchiolitis	9 (60.0)	9 (60.0)	
Bronchopneumonia	6 (40.0)	6 (40.0)	
Past history of prematurity	2 (13.3)	3 (20.0)	
Clinically significant illness	8 (53.3)	9 (60.0)	
Underlying chronic illness	3 (20)*	3 (20)†	
Home oxygen/respiratory support	2 (13.3)	1 (6.7)	
<b>Parameter at enrolment</b>			
Respiratory rate (breath/min)	42 (38–48)	45 (35–48)	0.819
Heart rate (beat/min)	130 (125–142)	130 (121–139)	0.803
Oxygen saturation (%)	97 (92–99)	98 (95–98)	0.641
Temperature	36.7 (36.5–37.0)	36.6 (36.4–36.9)	0.167
<b>mRDAI score</b>			
Crackles	5 (4–5)	4 (2–4)	
Wheeze	5 (4–5)	4 (4–5)	
Retraction	4 (3–5)	4 (3–5)	
Total mRDAI score	11 (10–12)	11 (8–12)	0.486
Wang clinical severity score	9 (7–9)	9 (7–9)	0.683
FLACC score	3 (0–6)	5 (0–6)	0.528
<b>Treatment of current illness</b>			
Bronchodilators	2 (13.3)	1 (6.7)	
Antibiotics	8 (53.3)	8 (53.3)	
Oxygen therapy at enrolment	6 (40)	6 (40)	
<b>Culture positive</b>			
Respiratory syncytial virus	2 (13.3)	1 (6.7)	
Other viral pathogens	2	1	
Bacterial pathogens	3	2	
Length of stay (n = 15)	4 (3–8)	5 (4–6)	0.683
Acute bronchiolitis (n = 9)	4 (3–7.5)	4 (3.5–5.5)	0.863
Bronchopneumonia (n = 6)	5 (3.5–9.3)	5.5 (4.75–7.8)	0.699

Underlying chronic illnesses include \*cystic fibrosis, neurodevelopmental delay, trisomy 21 and chronic lung disease; †cerebral palsy, neurodevelopmental delay, trisomy 21 and obstructive sleep apnoea syndrome. CPT: chest physiotherapy; FLACC: Face, Leg, Activity, Cry, Consolability scale; IQR: interquartile range; mRDAI: modified Respiratory Distress Assessment Instrument

Table III. Changes in clinical characteristics of patients before ( $T_0$ ) and after intervention ( $T_{120}$ ).

Variable	Manual CPT (n = 45)				Mechanical CPT (n = 42)				p-value <sup>†</sup>
	$T_0$	$T_{120}$	$T_{120} - T_0$	p-value*	$T_0$	$T_{120}$	$T_{120} - T_0$	p-value*	
mRDAI score	10.7 ± 2.2	7.7 ± 2.5	-3.0 ± 2.1	0.0001	10.3 ± 2.1	6.5 ± 2.4	-3.6 ± 2.1	0.0001	0.144
Wheezing	2.9 ± 1.7	2.0 ± 1.6	-1.0 ± 1.5	0.0001	3.1 ± 1.8	1.6 ± 1.7	-1.6 ± 1.8	0.0001	0.096
Crackles	4.0 ± 1.1	2.9 ± 1.4	-1.1 ± 1.3	0.0001	3.6 ± 1.3	2.7 ± 1.4	-0.9 ± 1.5	0.0001	0.584
Retractions	3.8 ± 0.9	2.9 ± 1.0	-0.9 ± 1.1	0.0001	3.6 ± 1.0	2.5 ± 1.1	-1.2 ± 1.0	0.0001	0.244
Respiratory rate	38.7 ± 7.7	35.1 ± 7.4	-3.6 ± 7.1	0.002	40.9 ± 7.3	34.0 ± 7.8	-7.0 ± 6.7	0.0001	0.024 <sup>‡</sup>
Heart rate	127.8 ± 13.7	127.1 ± 14.1	-0.6 ± 12.1	0.722	131.2 ± 15.3	128.0 ± 14.1	-3.2 ± 14.9	0.171	0.378
SpO <sub>2</sub>	96.6 ± 3.2	97.2 ± 3.0	0.6 ± 2.4	0.123	96.6 ± 2.6	97.5 ± 2.1	0.9 ± 2.2	0.013	0.509
Temperature	36.6 ± 0.4	36.7 ± 0.6	0.0 ± 0.6	0.757	36.6 ± 0.4	36.7 ± 0.5	0.1 ± 0.3	0.256	0.747

Data presented as mean ± standard deviation, unless otherwise stated. \*Paired t-test comparing pre- ( $T_0$ ) and post ( $T_{120}$ ) intervention. †Student's t-test comparing manual CPT and mechanical CPT. ‡p < 0.05 was considered statistically significant. mRDAI: modified Respiratory Distress Assessment Instrument; SpO<sub>2</sub>: oxygen saturation via pulse oximetry

**Table IV. FLACC score before ( $T_0$ ), during ( $T_{CPT}$ ) and after ( $T_{120}$ ) intervention.**

CPT group	FLACC score*						
	$T_0$	$T_{CPT}$	$T_{120}$	$T_{CPT} - T_0$	p-value	$T_{120} - T_{CPT}$	p-value
Manual (n = 45)	3.56 ± 2.74	3.80 ± 2.4	0.84 ± 1.24	0.24 ± 1.2	0.175	-2.96 ± 2.55	0.0001
Mechanical (n = 42)	3.29 ± 2.61	3.71 ± 2.13	1.0 ± 1.47	0.43 ± 1.43	0.06	-2.71 ± 2.8	0.0001
p-value	0.64	0.86	0.59	0.50		0.66	

Data presented as mean ± standard deviation, unless otherwise stated. \*FLACC score between 0 and 10, with 0 = relaxed and comfortable; 1–3 = mild distress; 4–6 = moderate pain; and 7–10 = severe distress. CPT: chest physiotherapy; FLACC: Face, Leg, Activity, Cry, Consolability scale

each group) at enrolment to five on Day 4. No adverse events such as bradycardia, desaturation, vomiting or rib fractures were observed during CPT. There was no difference in the median duration of hospitalisation between the manual and mechanical groups (4.4 days vs. 4.88 days,  $p = 0.654$ ).

## DISCUSSION

In this study, we observed significant improvements in RR, oxygen saturation and mRDAI in patients who underwent either manual or mechanical CPT. Children using mechanical CPT had greater reduction in their RR and more improvement in oxygen saturation after intervention. No side effects were noted, and improvements in the FLACC score were observed after both interventions.

The baseline demographics in both groups were comparable for age, weight and severity of respiratory distress. While there was a male preponderance in the mechanical group, this is unlikely to have influenced the results. The median mRDAI score of 11 at enrolment indicated moderately severe respiratory distress and a need for CPT. The efficacy of the LEGA-Kid machine was comparable to that of manual CPT performed by an experienced chest physiotherapist. The main drawbacks to performing effective manual CPT include the need for trained personnel, variability in techniques and skills among personnel, time constraints and familial burden. The combination of percussion and vibrations provided by the LEGA-Kid machine could mimic manual CPT, which helps to dislodge loosened secretions. Maximal intrathoracic pressure was achieved when the percussor was pressed firmly and at right angles to the chest wall to make the voice quiver. The percussor was shown to maintain a higher and more constant intrathoracic pressure, comparable to the performance of three physiotherapists.<sup>(16)</sup>

Our results differ from those of previous studies that compared CPT versus no CPT in children less than two years of age with severe bronchiolitis; no significant improvement in clinical severity was reported in children who received CPT compared to those who did not.<sup>(17-19)</sup> These studies, however, did not incorporate prior nebulisation with hypertonic saline. Of the 12 randomised controlled trials reviewed in the 2016 Cochrane review,<sup>(9)</sup> only one study<sup>(8)</sup> included the use of nebulised 3% saline with albuterol with or without CPT in patients with moderately severe respiratory syncytial virus bronchiolitis. An immediate reduction in wheezing and retractions was observed in patients who received CPT, with no impact on duration of hospitalisation.<sup>(8)</sup>

Despite the heterogeneity of the patients in our study, all of them retained airway secretions, which contributed to their respiratory distress. Nebulised hypertonic saline induces an

osmotic flow of water in the inspissated mucous, thus reducing the viscosity of the mucous and oedema in the submucosal tissue.<sup>(20)</sup> Subsequent CPT assists in the clearance of the loosened secretions, resulting in improved tidal volume, a more effective cough and further removal of secretions, which immediately helps to make breathing easier for the young child. This sequence of events explains the improvement in RR and better scores for the three components of mRDAI at  $T_{120}$  as compared to those at  $T_0$ . Our combined strategy of nebulised hypertonic saline followed by CPT differs from that of studies that investigated the efficacy of nebulised hypertonic saline alone in the management of LRTI.<sup>(21-23)</sup>

Both the manual and mechanical CPT procedures used in our study were well tolerated by the patients, as measured by the FLACC scale. Our patients' acceptance of CPT could be partially explained by the distraction methods used. The FLACC score was further reduced at  $T_{120}$ , indicating an improvement in overall condition after intervention. No adverse events were reported. The median length of stay in patients with acute bronchiolitis and bronchopneumonia was similar to those reported in the literature.<sup>(17,24)</sup>

The strengths of our study lie in the standardisation of CPT methods with the use of one trained physiotherapist to perform both the mechanical and manual CPT, and two physicians throughout the study – one to evaluate safety and tolerability, and the other to assess the efficacy of CPT. Furthermore, the latter physician was blinded to the method of CPT. However, our study was not without its limitations. First, it was limited by the lack of a control group with no CPT, which may have affected the interpretation of our results. We were unable to have a control group, as our hospital's policy requires children with LRTI who are aged below two years to be managed with regular CPT when needed. Second, we did not conduct subsequent evaluations beyond two hours to determine whether the improvements observed were sustained; nonetheless, no child in the study deteriorated during or after CPT. Third, our study also did not set out to investigate the ease of the percussor versus conventional physiotherapy. Finally, our sample size was small but sufficient for a pilot study.

In summary, as an adjunct treatment for LRTI, CPT is safe and beneficial for mucociliary clearance and improves moderately severe respiratory distress. The LEGA-Kid percussor is better than manual CPT performed by experienced personnel. However, further studies are needed to confirm these preliminary results.

## ACKNOWLEDGEMENTS

We thank the following individuals for their contribution to the study: A/Prof Thavagnanam Surendran, A/Prof de Bruyne Jessie

Anne, A/Prof Gan Chin Seng, Dr Chuah Soo Lin and Dr Abdul Aziz Fadzlina; the medical and nursing staff of Paediatric Respiratory Ward, Department of Paediatrics; and Manjit Singh Anmol Kaur and Haron Norazah, Physiotherapy Division, Department of Rehabilitation Medicine, UMMC, Kuala Lumpur, Malaysia. The study was funded by PlaTCOM Ventures Sdn Bhd via a grant to Formedic Technology (grant PlaTCOM-HIP-2015/013). Neither PlaTCOM Ventures Sdn Bhd nor Formedic Technology had any role in the study design, data collection and analysis, manuscript preparation or the decision to publish the manuscript.

## REFERENCES

1. Leowski J. Mortality from acute respiratory infections in children under 5 years of age: global estimates. *World Health Stat Q* 1986; 39:138-44.
2. Griscom NT, Wohl ME, Kirkpatrick JA Jr. Lower respiratory infections: how infants differ from adults. *Radiol Clin North Am* 1978; 16:367-87.
3. Wallis C, Prasad A. Who needs chest physiotherapy? Moving from anecdote to evidence. *Arch Dis Child* 1999; 80:393-7.
4. Lukrafka JL, Fuchs SC, Fischer GB, et al. Chest physiotherapy in paediatric patients hospitalised with community-acquired pneumonia: a randomised clinical trial. *Arch Dis Child* 2012; 97:967-71.
5. Rochat I, Leis P, Bouchardy M, et al. Chest physiotherapy using passive expiratory techniques does not reduce bronchiolitis severity: a randomised controlled trial. *Eur J Pediatr* 2012; 171:457-62.
6. Postiaux G, Louis J, Labasse HC, et al. Evaluation of an alternative chest physiotherapy method in infants with respiratory syncytial virus bronchiolitis. *Respir Care* 2011; 56:989-94.
7. Remondini R, Santos AZ, Castro G, Prado C, Silva Filho LV. Comparative analysis of the effects of two chest physical therapy interventions in patients with bronchiolitis during hospitalization period. *Einstein (Sao Paulo)* 2014; 12:452-8.
8. Gomes EL, Postiaux G, Medeiros DR, et al. Chest physical therapy is effective in reducing the clinical score in bronchiolitis: randomized controlled trial. *Rev Bras Fisioter* 2012; 16:241-7.
9. Roqué i Figuls M, Giné-Garriga M, Granados Rugeles C, Perrotta C, Vilaró J. Chest physiotherapy for acute bronchiolitis in paediatric patients between 0 and 24 months old. *Cochrane Database Syst Rev* 2016; 2:CD004873.
10. Punithavathi N, Ong LM, Irfhan Ali HA, et al. A pilot randomized control cross over study evaluating the effectiveness and safety of mechanical percussor compared with conventional chest physiotherapy in adults with productive cough. *Med J Malaysia* 2014; 69:16-20.
11. Urbaniak GC, Plous S. Research Randomizer (version 4.0): computer software, 2015 (updated January 20, 2017). Available at: <http://www.randomizer.org/>. Accessed [DATE]
12. Lowell DI, Lister G, Von Koss H, McCarthy P. Wheezing in infants: the response to epinephrine. *Pediatrics* 1987; 79:939-45.
13. Wang EE, Milner RA, Navas L, Maj H. Observer agreement for respiratory signs and oximetry in infants hospitalized with lower respiratory infections. *Am Rev Respir Dis* 1992; 145:106-9.
14. Chin HJ, Seng QB. Reliability and validity of the respiratory score in the assessment of acute bronchiolitis. *Malays J Med Sci* 2004; 11:34-40.
15. Manworren RC, Hynan LS. Clinical validation of FLACC: preverbal patient pain scale. *Pediatr Nurs* 2003; 29:140-6.
16. Flower KA, Eden RI, Lomax L, Mann NM, Burgess J. New mechanical aid to physiotherapy in cystic fibrosis. *Br Med J* 1979; 2:630-1.
17. Webb MSC, Martin JA, Carlidge PHT, Ng YK, Wright NA. Chest physiotherapy in acute bronchiolitis. *Arch Dis Child* 1985; 60:1078-9.
18. Nicholas KJ, Dhouieb MO, Marshall TG, Edmunds AT, Grant MB. An evaluation of chest physiotherapy in the management of acute bronchiolitis: changing clinical practice. *Physiotherapy* 1999; 85:669-74.
19. Bohé L, Ferrero M, Cuestas E, Polliotto L, Genoff M. [Indications of conventional chest physiotherapy in acute bronchiolitis]. *Medicina (B Aires)* 2003; 64:198-200. Spanish.
20. Mandelberg A, Amirav I. Hypertonic saline or high volume normal saline for viral bronchiolitis: mechanisms and rationale. *Pediatr Pulmonol* 2010; 45:36-40.
21. Köse S, Şehriyaroglu A, Esen F, et al. Comparing the efficacy of 7%, 3% and 0.9% saline in moderate to severe bronchiolitis in infants. *Balkan Med J* 2016; 33:193-7.
22. Zhang L, Mendoza-Sassi RA, Klassen TP, Wainwright C. Nebulized hypertonic saline for acute bronchiolitis: a systematic review. *Pediatrics* 2015; 136:687-701.
23. Baron J, El-Chaar G. Hypertonic saline for the treatment of bronchiolitis in infants and young children: a critical review of the literature. *J Pediatr Pharmacol Ther* 2016; 21:7-26.
24. Paludo C, Zhang L, Lincho CS, et al. Chest physical therapy for children hospitalised with acute pneumonia: a randomised controlled trial. *Thorax* 2008; 63:791-4.