

# Anaesthesia and intensive care medicine in the COVID-19 pandemic

Jolin Wong<sup>1</sup>, MMed, FANZCA, Shin Yi Ng<sup>1</sup>, MMed, Meng Huat Goh<sup>1</sup>, MMed, FANZCA, Kian Hian Tan<sup>1</sup>, FANZCA, FPPMANZCA, Hui Zhong Chai<sup>2</sup>, MBBS, MRCP, Ken Junyang Goh<sup>2</sup>, MD, MRCP, Chai Rick Soh<sup>1</sup>, FANZCA, FCICM

## INTRODUCTION

The COVID-19 pandemic has affected millions and resulted in countless deaths worldwide. Healthcare systems have been overwhelmed in many countries, including China, Italy,<sup>(1)</sup> Spain, the United Kingdom<sup>(2)</sup> and the United States. This is particularly so for critical care medicine, where infrastructure, drugs, equipment and skilled staff may be in short supply.

Singapore was one of the first countries to be affected by the early wave of infections connected to Wuhan, China. All restructured hospitals and some private hospitals have played major roles in Singapore's pandemic response so far, including screening for COVID-19, caring for afflicted patients, boosting manpower at the National Centre for Infectious Diseases and sending teams into the community to screen for the virus. Anaesthesiologists and intensive care physicians have been an integral part of Singapore's healthcare response. This article describes the potential and real contributions by anaesthesiologists and intensivists in the pandemic response efforts.

## SAFETY IN THE OPERATING THEATRE

An anaesthesiologist's routine work involves multiple aerosol-generating procedures, including intubation, extubation and suctioning of the airway. Experience during the severe acute respiratory syndrome outbreak in 2003 has shown that endotracheal intubation carries the highest risk of viral transmission to the healthcare worker.<sup>(3)</sup>

The Division of Anaesthesiology in Singapore General Hospital (SGH) was quick to respond to the initial outbreak of the novel respiratory pathogen by preparing a set of guidelines for anaesthesia and perioperative care of patients to minimise potential transmission of infection.<sup>(4-6)</sup> Traditionally considered leaders in patient safety, we also created new processes and systems in our operating theatres (OTs), designating an isolation OT that was physically separated from the main OT complex. With strong support from the OT nursing staff, *in situ* simulation sessions were performed early to ensure that all team members were prepared for the eventuality of a COVID-19 patient requiring surgery. COVID-19 coordinators were present at all operations performed on suspected or confirmed COVID-19 patients, providing guidance to all staff, including surgeons, nurses and healthcare attendants, to ensure a safe environment for both patients and healthcare workers. Since then, anaesthesiologists the world over have come up with many different guidelines for anaesthetic and airway management for COVID-19 patients.

Similar guidelines have been created by various medical specialties across Singapore.<sup>(7-11)</sup>

## SAFETY IN EMERGENCY DEPARTMENT AND PRIMARY CARE

As the outbreak spread globally into a pandemic, Singapore experienced a second wave of imported infections, with a rise in the number of COVID-19-related attendances at the emergency department. Armed with our airway expertise, the division sent teams to the emergency department and fever screening areas to assist in obtaining nasopharyngeal and throat swabs for patients suspected to have COVID-19. Through this experience, innovative minds came together and created the SG-Swab Assurance for Everyone booths to protect doctors, as they performed swabs.<sup>(12)</sup> Further work was done to create protective boxes and screens for intubation to reduce droplet spread of the virus during airway manipulation.<sup>(13)</sup> Acceleration of these developments was possible owing to the availability of emergency funds that could be rapidly disbursed for COVID-19-related projects. Furthermore, throughout these efforts, philanthropy was encouraged to support safety and innovation in healthcare.

## Creating surge capacity for intensive care

At the time of writing, evidence indicates that 5%–9% of all COVID-19 patients require intensive care.<sup>(14,15)</sup> As the number of infected patients rises in Singapore, it is imperative to create critical care surge capacity.<sup>(16)</sup> The team from the Department of Surgical Intensive Care, SGH, comprising anaesthesiologist-intensivists, was involved in preparations for a surge in critically ill patients as part of coordinated efforts by the national intensive care unit (ICU) committee. Significant work was put in to map the expansion of our isolation ICU, including the conversion of general ward beds to critical care beds. On top of our usual clinical work, ICU teams, comprising a consultant intensivist, a senior anaesthesiology resident and medical officers, have been assembled to care for critically ill COVID-19 patients. Our nurses in the OT are well placed for rapid training to serve as ICU nurses during a surge.

There is significant overlap between care of the critically ill patient and care of the anaesthetised patient undergoing surgery. All anaesthesiologists have gained valuable experience in the ICU as part of their specialty training and may be called upon to lead ICU teams in a surge. The skills that anaesthesiologists have developed will prove essential in the ICU. For example, we are

<sup>1</sup>Division of Anaesthesiology, <sup>2</sup>Department of Respiratory and Critical Care Medicine, Singapore General Hospital, Singapore

**Correspondence:** Dr Jolin Wong, Consultant, Division of Anaesthesiology, Singapore General Hospital, Outram Road, Singapore 169608. [jolin.wong@singhealth.com.sg](mailto:jolin.wong@singhealth.com.sg)

experts in airway management and vascular access, especially for invasive lines. Ventilator management is second nature, and haemodynamic monitoring is *de rigueur*. Skills in certain procedures, such as turning a patient prone, have turned out to be an unexpected asset during this pandemic, where many of the patients suffering from acute respiratory distress syndrome require prone positioning as part of respiratory management. These skills allow anaesthesiologists to function well as leaders of procedural teams such as airway or code teams, which may then be supported by clinicians and nurses who are not domain experts in these fields.<sup>(17)</sup>

The OT is a place that can be quickly converted into a temporary ICU should the need arise.<sup>(18)</sup> Both individual OTs and post-anaesthetic care unit (PACU) cubicles are readily suited to monitor critically ill patients, and come with oxygen source, wall-mounted suction and multiple power points. However, the conversion of such locations into an ICU is not without significant effort, as engineering modifications may be required to enhance the capabilities of the OT complex. For instance, taps and drainage outlets are required for haemodialysis, and these are not part of the usual features of an OT. Concessions may have to be made for the level of care depending on the capabilities of the location. For instance, although OTs normally have outlets for medical air, a PACU does not. As a result, the ventilators used in the PACU must be able to use entrained atmospheric air. In an OT that has no tap or drainage, continuous renal replacement therapy must be used instead of intermittent haemodialysis.

In a pandemic, vast quantities of supplies and drugs need to be prepared to cope with the surge in critically ill patients. Many physicians in acute medicine have attempted to think out of the box to enhance the utilisation of existing equipment to care for larger numbers of patients.<sup>(19)</sup> Assessments on the safety of such modifications, however, fall back on basic understanding of physiology and ventilatory management; as such, these suggestions remain outside the scope of conventional critical care practice.<sup>(20)</sup> The intimate knowledge that anaesthesiologists possess about their equipment has been invaluable in engineering creative methods to substitute existing consumables for those that may run short. For example, with the help of our anaesthetic technicians, we used circle circuits to replace single-limb breathing circuits in the transport ventilators. We also reduced dependency on disposable items, preferentially choosing to use reusable alternatives. Furthermore, we worked closely with the sterile supplies unit to re-process certain single-use items to cope with the large numbers required. Apart from medical equipment, anaesthesiologists are also involved in the conservation of personal protective equipment, which is of vital importance in protecting healthcare workers during an infectious disease outbreak.<sup>(21)</sup>

Anaesthesiologists' knowledge of pharmacology has been extremely useful in brainstorming alternative drugs and drug-conservation strategies to cope with the increase in demand for drugs used in the ICU. Drugs such as thiopentone and pancuronium, which had somewhat fallen out of favour in recent years, have reappeared in the anaesthesiologists'

armamentarium to spare the use of propofol and atracurium in the OT. Anaesthesiologists are also now relying on central neuraxial and peripheral nerve blocks for similar reasons. Finally, our knowledge of gases and inhalational agents has led us to consider and examine the hospital's vacuum-insulated evaporator and oxygen supply for their capacity to cope with a surge in demand for mechanical ventilation.<sup>(22)</sup>

With the need to maintain safe distancing, we have come to embrace the use of technology for the continuation of education, peer-reviewed learning and administrative meetings in the department. Surgical intensivists have always been involved in training surgical residents to care for critically ill patients. In the current climate, as all available manpower is mobilised in the ICU surge, we have also ramped up education for our surgical colleagues to refresh their knowledge in intensive care. This enhances capability on the ground to take care of more patients. In collaboration with our colleagues in Respiratory and Critical Care Medicine, a set of lectures and ventilator demonstrations have been recorded as part of an online ICU refresher course.<sup>(23)</sup> This enables all physicians, surgeons and junior doctors to be quickly trained as non-intensivist members of the ICU team. Finally, as the most frequent users of powered air-purifying respirators, anaesthesiologists have also taken on the role of coaches to operating team members who may be unfamiliar with their personal protective equipment.

The COVID-19 pandemic threatens to last for a protracted amount of time. Although the pandemic presently remains a global crisis, it may give rise to opportunities to change our practice for the better and, at the same time, encourage collaboration and medical innovation to benefit our patients.<sup>(24,25)</sup> To emerge victorious, we will require resilience. As leaders of a medical team, we should vigilantly look out for evidence of burnout among our fellow healthcare workers and provide one another with the necessary support systems to get through this difficult time together.<sup>(26)</sup>

## REFERENCES

1. Armocida B, Formenti B, Ussai S, Palestra F, Missoni E. The Italian health system and the COVID-19 challenge. *Lancet Public Health* 2020; 5:e253.
2. Horton R. Offline: COVID-19 and the NHS—"a national scandal". *Lancet* 2020; 395:1022.
3. Tran K, Cimon K, Severn M, Pessoa-Silva CL, Conly J. Aerosol generating procedures and risk of transmission of acuterespiratory infections to healthcare workers: a systematic review. *PLoS One* 2012; 7:e35797.
4. Wong J, Goh QY, Tan Z, et al. Preparing for a COVID-19 pandemic: a review of operating room outbreak response measures in a large tertiary hospital in Singapore. *Can J Anaesth* 2020; 67:732-45.
5. Lie SA, Wong SW, Wong LT, Wong TGL, Chong SY. Practical considerations for performing regional anesthesia: lessons learned from the COVID-19 pandemic. *Can J Anaesth* 2020; 67:885-92.
6. Ong S, Tan TK. Practical considerations in the anaesthetic management of patients during a COVID-19 epidemic. *Anaesthesia* 2020; 75:823-4.
7. Chew MH, Koh FH, Ng KH. A call to arms: a perspective of safe general surgery in Singapore during the COVID-19 pandemic. *Singapore Med J* 2020; 61:378-80.
8. Ang TL, Li JW, Vu CK, et al. Chapter of Gastroenterologists professional guidance on risk mitigation for gastrointestinal endoscopy during COVID-19 pandemic in Singapore. *Singapore Med J* 2020; 61:345-9.
9. Cherian R, Poh KK. At the 'heart' of the COVID-19 outbreak: early cardiac implications and mitigation strategies. *Singapore Med J* 2020; 61:373-4.
10. Kok SXS, Shah MT, Cheong WK, et al. Dealing with COVID-19: initial perspectives of a small radiology department. *Singapore Med J* 2020; 61:375-7.

11. Tay K, Kamarul T, Lok WY, et al. COVID-19 in Singapore and Malaysia: rising to the challenges of orthopaedic practice in an evolving pandemic. *Malays Orthop J* 2020; 14:7-15.
12. Teo J. Extra protection for healthcare workers doing large-scale Covid-19 tests. *The Straits Times* 2020 Apr 17. Available at: <https://www.straitstimes.com/singapore/extra-protection-for-healthcare-workers-doing-large-scale-covid-19-tests>. Accessed April 20, 2020.
13. Au Yong PS, Chen X. Reducing droplet spread during airway manipulation: lessons from the COVID-19 pandemic in Singapore. *Br J Anaesth* 2020; 125:e176-8.
14. Grasselli G, Zangrillo A, Zanella A, et al. Baseline characteristics and outcomes of 1591 patients infected with SARS-CoV-2 admitted to ICUs of the Lombardy region, Italy. *JAMA* 2020; 323:1574-81.
15. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirusdisease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention. *JAMA* 2020; 323:1239-42.
16. Grasselli G, Pesenti A, Cecconi M. Critical care utilization for the COVID-19 outbreak in Lombardy, Italy: early experience and forecast during an emergency response. *JAMA* 2020; 323:1545-6.
17. Zarzaur BL, Stahl CC, Greenberg JA, Savage SA, Minter RM. Blueprint for restructuring a department of surgery in concert with the health care system during a pandemic: the University of Wisconsin experience. *JAMA Surg* 2020; 155:628-35.
18. Phua J, Weng L, Ling L, et al. Intensive care management of coronavirus disease 2019 (COVID-19): challenges and recommendations. *Lancet Respir Med* 2020; 8:506-17.
19. Neyman G, Irvin CB. A single ventilator for multiple simulated patients to meet disaster surge. *Acad Emerg Med* 2006; 13:1246-9.
20. American Society of Anesthesiologists. Joint statement on multiple patients per ventilator. Available at: <https://www.asahq.org/about-asa/newsroom/news-releases/2020/03/joint-statement-on-multiple-patients-per-ventilator>. Accessed June 1, 2020.
21. Livingston E, Desai A, Berkwits M. Sourcing personal protective equipment during the COVID-19 pandemic. *JAMA* 2020; 323:1912-4.
22. Moonesinghe R, Menzies-Gow A, Eggleton A. Estates and Facilities Alert: Use of high flow oxygen therapy devices (including wall CPAP and high flow face mask or nasal oxygen) during the Coronavirus epidemic – urgent patient safety notice; immediate attention required. Department of Health, United Kingdom. 31 March 2020. Available at: [https://www.cas.mhra.gov.uk/ViewandAcknowledgment/ViewAttachment.aspx?Attachment\\_id=103492](https://www.cas.mhra.gov.uk/ViewandAcknowledgment/ViewAttachment.aspx?Attachment_id=103492). Accessed June 2, 2020.
23. Chai HZ, Teo CWS, Ng LS, et al. Rapid training of non-intensivists using an online critical care course during COVID-19. *Ann Acad Med Singap* 2021; 50:503-7.
24. Hsu LY, Chia PY, Vasoo S. A midpoint perspective on the COVID-19 pandemic. *Singapore Med J* 2020; 61:381-3.
25. Lum LHW, Tambyah PA. Outbreak of COVID-19 - An urgent need for good science to silence our fears? *Singapore Med J* 2020; 61:55-7.
26. Chew QH, Wei KC, Vasoo S, Chua HC, Sim K. Narrative synthesis of psychological and coping responses towards emerging infectious disease outbreaks in the general population: practical considerations for the COVID-19 pandemic. *Singapore Med J* 2020; 61:350-6.