

# Reference range for ADAMTS13 antigen, activity and anti-ADAMTS13 antibody in the healthy adult Singapore population

Allison Ching Yee Tso<sup>1</sup>, MRCP, FRCPath, Christina Lai Lin Sum<sup>2</sup>, MSc, Kiat Hoe Ong<sup>1</sup>, MRCP, FRCPA

**INTRODUCTION** ADAMTS13 (a disintegrin-like and metalloproteinase with a thrombospondin Type 1 motif, member 13) plays a fundamental role in the regulation of haemostasis and thrombosis. Its deficiency leads to an accumulation of ultra-large von Willebrand multimers, inducing spontaneous platelet aggregation, thrombosis in the microvasculature, and thrombotic thrombocytopenic purpura (TTP), a condition with 90% mortality when left untreated. Prompt quantification of ADAMTS13 antigen, activity and autoantibody plays a crucial role in the diagnosis and management of TTP and can help differentiate it from other thrombotic microangiopathies (TMAs). Reference ranges for ADAMTS13 are generally derived from Caucasian patients. Given that polymorphism in the ADAMTS13 gene can be associated with variable ADAMTS13 levels, we aimed to establish the first reference range in Singapore and provide a crucial laboratory test for institutions here and elsewhere.

**METHODS** 150 healthy voluntary donors (75 men, 75 women) aged 21–60 years, with an ethnic mix mirroring Singapore's population profile, were recruited. ADAMTS13 antigen, activity and autoantibody levels were measured using the fluorescence resonance energy transfer-vWF73 and enzyme-linked immunosorbent assay methodologies.

**RESULTS** Levels (activity 0.65–1.79 IU/mL, antigen 0.36–1.17 IU/mL, autoantibody 1.4–12.5 U/mL) were not statistically different between the genders and various age groups.

**CONCLUSION** TTP and TMAs are encountered in a wide range of specialties. The availability of new assays in Singapore will aid clinicians in the timely management of these conditions. Standardising reference ranges established for Singapore against World Health Organization standards allows harmonisation of measurements between laboratories and for future research collaborations.

*Keywords:* ADAMTS13, microangiopathic haemolytic anaemia, thrombotic thrombocytopenic purpura

## INTRODUCTION

ADAMTS13 (a disintegrin-like and metalloproteinase with a thrombospondin Type 1 motif, member 13) is an essential protease that regulates the size of ultra-large von Willebrand factor (UL-vWF) by cleavage at the Tyr1605-Met1606 bond in the A2 domain.<sup>(1)</sup> Under conditions of shear stress in the presence of ADAMTS13 deficiency, UL-vWF multimers accumulate, which, in turn, induce spontaneous platelet aggregation and thrombi formation in the microvasculature, especially in vital organs, leading to thrombotic thrombocytopenic purpura (TTP).

Up to 20% of patients with TTP present with neurological complications,<sup>(2)</sup> 11% of patients have acute and/or chronic renal failure,<sup>(3)</sup> and patients can experience sudden cardiac death secondary to myocardial infarction, heart failure and arrhythmias.<sup>(4)</sup> Without treatment, TTP is associated with a mortality rate of about 90%, with a majority of deaths occurring within the first 24 hours of presentation.

TTP is a rare disorder, with a reported annual incidence of six per million population.<sup>(5)</sup> Congenital TTP (or Upshaw-Schulman syndrome) is due to an inherited deficiency of ADAMTS13, while acquired TTP results from a deficiency of ADAMTS13 secondary to anti-ADAMTS13 antibodies. The ability to quantify ADAMTS13 activity and autoantibody levels

is critical for differentiating inherited TTP from acquired TTP. It would also help to differentiate TTP from other thrombotic microangiopathies such as haemolytic uraemic syndrome (HUS), atypical HUS, disseminated intravascular coagulation, HELLP (haemolysis, elevated liver enzymes, low platelet) syndrome/pre-eclampsia, malignant hypertension,<sup>(6)</sup> Kasabach-Merritt syndrome, paroxysmal nocturnal haemoglobinuria and catastrophic antiphospholipid syndrome. The absence of ADAMTS13 autoantibodies in a patient with < 10% ADAMTS13 activity differentiates congenital TTP from acquired TTP, as 94%–97% of patients suffer from idiopathic TTP and have severe ADAMTS13 deficiency but will have circulating anti-ADAMTS13 antibodies in the plasma.<sup>(7)</sup> This undoubtedly affects the modalities of patient management, as congenital TTP is treated with regular infusion of plasma and, potentially, the novel therapy BAX 930/SHP655, a recombinant ADAMTS13,<sup>(8,9)</sup> which is still in clinical trials. On the other hand, the treatment for acquired TTP requires plasma exchange, corticosteroids and/or rituximab (anti-CD20 monoclonal antibody).

Up to 75%–90% of patients with TTP would have very low ADAMTS13 activity level < 10%, a reading that has 90% sensitivity and specificity. In two more recent studies, ADAMTS13 activity cut-off < 10% was found in more than 97.5%<sup>(10)</sup> and 100%<sup>(11)</sup> of patients with TTP. The median ADAMTS13 activity of patients

<sup>1</sup>Department of Haematology, <sup>2</sup>Department of Laboratory Medicine, Tan Tock Seng Hospital, Singapore

**Correspondence:** Dr Allison Ching Yee Tso, Consultant, Department of Haematology, Tan Tock Seng Hospital, 11 Jalan Tan Tock Seng, Singapore 308433. [allison\\_cy\\_tso@ttsh.com.sg](mailto:allison_cy_tso@ttsh.com.sg)

with thrombotic microangiopathies in the United Kingdom TTP registry, on the other hand, ranges from 56% to 66%.<sup>(5)</sup> However, it should be stated that up to 14% of patients with HUS<sup>(12)</sup> and up to 10% of patients with severe Child-Pugh C liver cirrhosis can have ADAMTS13 activity level < 10%.<sup>(13)</sup>

ADAMTS13 activity and autoantibody levels can provide prognostic information in acute TTP. Patients with activity level < 10% are more likely to respond to plasma exchange, have higher remission rate (> 80% vs. 20%–75% in three different studies<sup>(14–16)</sup>) and lower mortality rate (16% vs. 45%<sup>(14)</sup>).<sup>(17)</sup> A prospective study conducted by Ferrari et al found the persistence of undetectable activity in remission to be predictive of relapse within 18 months, with a positive predictive value of 38.5%. In contrast, the negative predictive value for relapse of measurable ADAMTS13 activity in remission was 94.7%.<sup>(18,19)</sup>

Despite appropriate treatment, the mortality of patients with TTP still ranges between 15% and 20%.<sup>(5)</sup> Around 30% (in certain studies, up to 60%<sup>(20)</sup>) of patients who achieve remission would also relapse.<sup>(20)</sup> Persistence of ADAMTS13 autoantibodies during remission predicts recurrence,<sup>(21,22)</sup> with a threefold increase in relapse rate compared with those without detectable autoantibodies.<sup>(21)</sup> However, the timing of relapse is unpredictable, as demonstrated in one study, where a wide range was noted (duration 0.3–9.5 years).<sup>(23)</sup> Hence, although the ADAMTS13 assay cannot inform clinicians about the timing to commence pre-emptive treatment, it would strongly support the need for closer monitoring than in patients without persistent autoantibodies and low activity during clinical remission. The persistence of low ADAMTS13 activity level and elevated autoantibodies during acute TTP treatment, on the other hand, would suggest the need to intensify plasma exchange and/or institute additional therapy, such as rituximab.<sup>(24)</sup>

There is no universally agreed normal or reference range for ADAMTS13 activity. Variations exist depending on the assay methodology, age, gender and genetic variants in different ethnic populations.<sup>(25–27)</sup> In general, normal ADAMTS13 activity for healthy adults ranges between 50% and 160%. For example, the Mayo Clinic, USA, accepts an activity level  $\geq$  70% to be normal.<sup>(28)</sup> However, in parts of Europe,<sup>(29)</sup> the lower limit of reference range for ADAMTS13 activity can be as low as 40%. Single-nucleotide polymorphisms, for example P.Ala732Val, can be associated with a decrease of over 20% in ADAMTS13 activity. The P.Pro475Ser polymorphism, which is restricted to the Eastern Asian population, is heterozygous in approximately 10% of the Japanese population and associated with a reduction in ADAMTS13 activity level.<sup>(30)</sup> Feys et al reported that normal Chinese donors have significantly lower ADAMTS13 antigen levels than Caucasian donors.<sup>(31)</sup> It was not recommended to adopt the manufacturer's ADAMTS13 reference ranges for our laboratory, as the manufacturer used a different fluorescence reader and wavelengths (BioTek FL  $\times$  800 TBI, BioTek Instruments Inc, Winooski, VT, USA), with excitation at 340 nm and emission at 450 nm. Our study, meanwhile, used Tecan Infinite F200 Pro microplate reader (Tecan Austria GmbH, Grodig, Austria), with excitation at 340 nm and emission at 465 nm. Together with the variation of ADAMTS13 among

different ethnic groups, there was a need to establish the first ADAMTS13 reference range for adults in Singapore.

## METHODS

This prospective study recruited 150 healthy adult volunteers in Singapore. Ethics approval from the domain specific review board and informed consent of all participants were obtained prior to the study.

Blood (2  $\times$  2.7 mL citrate tube and 1  $\times$  3 mL EDTA [ethylenediaminetetraacetic acid] tubes) samples were collected from each volunteer. The equipment consisted of the Tecan Infinite F200 Pro microplate reader for fluorescent resonance energy transfer [FRET]-vWF73 and enzyme-linked immunosorbent assay [ELISA], Technozym® Fluorogenic ELISA kit, Technozym® ADAMTS-13 INH ELISA kit and the World Health Organization (WHO) First International Standard ADAMTS13 plasma (NIBSC code 12/252). Platelet-poor citrated plasma was prepared by centrifugation; the ADAMTS13 antigen and activity were measured using the FRET methodology, while human autoantibodies (immunoglobulin G [IgG]) in serum or plasma against ADAMTS13 were measured using ELISA. Full blood count was performed using the EDTA specimens. Platelet count was checked to be within the local reference interval.

For antigen and activity measurements, patients' samples, controls and calibrators were incubated with monoclonal anti-ADAMTS13 antibody coated on the wells of a microplate. The ADAMTS13 present was bound to the antibody. After washing, a fluorogenic substrate vWF-fragment (FRETS-vWF73) was added for activity measurement. Fluorescence generated by the cleavage of fluorogenic substrate was measured kinetically at 360 nm/460 nm for 15 minutes using the Tecan Infinite Series microplate reader. After measuring the ADAMTS13 activity, the microplate was washed. The conjugate, which contained anti-ADAMTS13 POX, was then added. This conjugate would then bind to the ADAMTS13 present. Excess conjugate was washed off and antigen substrate was added for colour development. Stop solution was then added to end the reaction after 15 minutes and the endpoint was read at 360 nm/460 nm. Patients' results were calculated from the reference curve generated. All calibrator values were calibrated against and traceable to the WHO First International Standard ADAMTS-13 plasma (12/252).

For autoantibody measurement, diluted samples, controls and calibrators were incubated with recombinant ADAMTS13 protease coated on the wells of a microplate. If autoantibodies were present in the sample, they would be bound to ADAMTS13 protease during incubation. After washing, antihuman IgG POX was added. This antibody binds to the ADAMTS13 autoantibodies. After incubation and washing, chromogenic substrate was added, which would react with the horseradish peroxidase of the antihuman IgG POX. After ten minutes of incubation, the reaction was stopped by adding stop solution. The intensity of colour was measured with Tecan Infinite Series microplate reader at 450 nm. A reference curve with point-to-point fit was set up to read the patient's result, where the X-axis represented concentration of ADAMTS13 IgG (U/mL) and Y-axis

represented extinction at 450 nm. Statistical software, including Microsoft Excel and Analyse-it, were used.

## RESULTS


The 150 (75 men, 75 non-pregnant women) healthy adult volunteers recruited were aged 21–60 years. The ethnic mix of donors mirrored the ethnic population profile of Singapore (115 Chinese, 20 Malay, 12 Indian, 3 Other). The donors were categorised into different age groups: 21–30 years; 31–40 years; 41–50 years; and 51–60 years. There was an equal number of male and female donors in each age group.

As the reference range for ADAMTS13 activity was 0.65–1.79 IU/mL, six of our 150 donors whose results exceeded the measurement range were excluded. Similarly, as the reference range for ADAMTS13 antigen was 0.36–1.17 IU/mL, two donors whose results exceeded the measurement range were excluded. Likewise, as the reference range for ADAMTS13 autoantibodies was 1.4–12.5 U/mL, one donor who was an outlier according to Dixon's rule was excluded. Samples with levels > 15 U/mL were considered as positive for autoantibodies. Anti-ADAMTS13 antibody below the detection limits was reported as < 2 U/mL.


There were no statistical differences, according to independent *t*-tests, for ADAMTS13 antigen (Fig. 1), activity (Fig. 1) and autoantibody (Fig. 2) levels between male and female volunteers and in the various age groups (Table I). However, as the number of recruits among the minority ethnicities was small (12 Indian donors and 20 Malay donors), we were unable to perform meaningful statistical analysis and draw conclusions with regard to the differences in ADAMTS13 antigen, activity and autoantibody levels between donors of various ethnicities.

## DISCUSSION

Historically, in Singapore, the diagnosis of TTP has been based on clinical features supported by laboratory findings of thrombocytopenia and peripheral blood film features of microangiopathic haemolytic anaemia. ADAMTS13 activity and autoantibodies, if obtained, are typically sent to the Mayo Clinic, USA, for analysis, but with long turnaround times of over two weeks. As the presenting features of TTP are often indistinguishable from those of other conditions, therapy for TTP, including plasmapheresis and corticosteroids that are not without risks, is often initiated as emergency treatment for suspected patients. With a turnaround time of  $\leq 2$  working days, the availability of this ADAMTS13 laboratory service in Singapore would have a significant impact on the clinical management of suspected TTP, providing clinicians the support to stop unnecessary treatment for patients for whom there is low clinical suspicion of TTP but who have nevertheless been commenced on plasma exchange, thereby reducing both risk to patients as well as healthcare expenditure. It can also be used to monitor the response to therapy and help predict relapses, especially for patients with chronic relapsing TTP. Furthermore, our reference range was standardised against WHO standard plasma, allowing reporting in international units (IU/mL), which is now encouraged by the WHO, as part of harmonisation of



**Fig. 1** Charts show ADAMTS13 activity and antigen levels according to (a) age group and (b) ethnicity. ADAMTS13: a disintegrin-like and metalloprotease with thrombospondin Type 1 motif, member 13



**Fig. 2** Charts show ADAMTS13 autoantibody levels according to (a) age group and (b) ethnicity. ADAMTS13: a disintegrin-like and metalloprotease with thrombospondin Type 1 motif, member 13

results across laboratories worldwide.

There are two types of ADAMTS13 inhibitors – neutralising antibodies (approximately two-thirds of patients) and non-neutralising antibodies (approximately one-third of patients). While neutralising antibodies block the substrate-cleaving activity, non-neutralising antibodies reduce the half-life of ADAMTS13

**Table I. ADAMTS13 antigen, activity and autoantibody levels according to gender, age group and ethnicity.**

Variable	Activity (IU/mL)			Antigen (IU/mL)			Autoantibody (U/mL)		
	No. of patients	Mean level	p-value*	No. of patients	Mean level	p-value*	No. of patients	Mean level	p-value*
<b>Total</b>	144	1.17	NA	150	0.77	NA	149	4.2	NA
<b>Gender</b>			0.23			0.46			0.97
Female	75	1.24		75	0.78		74	4.23	
Male	75	1.17		75	0.75		75	4.21	
<b>Age group (yr)</b>									
21–30	38	1.29	Ref	38	0.84	Ref	38	3.72	Ref
31–40	39	1.16	0.09	39	0.77	0.14	39	3.83	0.84
41–50	39	1.19	0.24	39	0.72	0.02	38	4.88	0.09
51–60	34	1.20	0.31	34	0.73	0.06	34	4.51	0.24
<b>Ethnicity</b>									
Chinese	115	1.22	Ref	115	0.78	Ref	115	4.33	Ref
Malay	20	1.19	0.72	20	0.74	0.48	20	4.18	0.83
Indian	12	1.13	0.38	12	0.71	0.26	11	2.8	0.08
Other	3	0.99	0.27	3	0.68	0.40	3	5.69	0.43

\*Independent t-test. ADAMTS13: a disintegrin-like and metalloprotease with thrombospondin type 1 motif, member 13; NA: not available

by enhancing antibody-mediated clearance. The advantage of our choice of assay (ELISA) to detect autoantibodies, over the traditional Bethesda assay utilised by some external reference laboratories, is the enhanced sensitivity that ELISA provides, as the Bethesda assay primarily detects only neutralising antibodies (seen in approximately two-thirds of patients) and cannot detect antibody titres < 0.5 BU (Bethesda units). In contrast, the ELISA methodology detects both types of inhibitors, thereby improving the sensitivity of the test as well as the diagnosis of TTP. One of the limitations of our assay was that samples with high concentrations of antibodies other than anti-ADAMTS13 antibodies might result in weak positive or borderline results.

Where the FRET assay is concerned, ADAMTS13 activity can be artificially low owing to high bilirubin (> 15 mg/dL) as well as haemolysed plasma containing free haemoglobin (> 2 g/dL). It is important to note the timing of the blood drawn in relation to whether the patient has already received plasma infusion or undergone plasma exchange, as the ADAMTS13 activity level would then be elevated. Activity can also be affected by high levels of endogenous vWF, the presence of heparin, fondaparinux, dabigatran or other direct thrombin inhibitors, and hyperlipidaemia.

As age can affect the ADAMTS13 activity by up to 3.9%–6.5%, the reference range established for healthy adults in Singapore may not be applicable to the paediatric population, despite our range being comparable to those used by paediatric centres in the USA.<sup>(32,33)</sup> Establishing a reference range of ADAMTS13 for the Singapore paediatric population is, therefore, warranted. Similarly, as genetic polymorphism can affect ADAMTS13 activity levels by up to 20%, we may not be able to confidently presume that the same cut-off values of ADAMTS13 activity in diseased states, as quoted in the guidelines, would apply to patients of Asian origin. Studying the pathological levels of ADAMTS13 in Singapore and Asian patients may inform practice guidelines and influence the clinical management of these patients.

## ACKNOWLEDGEMENTS

The authors thank the National Healthcare Group, Singapore, for its generous and unconditional support of this study through the awards of the Small Innovation Grant and Pitch-For-Fund Grant in 2015 and 2016, respectively.

## REFERENCES

- Xiang Y, de Groot R, Crawley JT, Lane DA. Mechanism of von Willebrand factor scissile bond cleavage by a disintegrin and metalloproteinase with a thrombospondin type 1 motif, member 13 (ADAMTS13). *Proc Natl Acad Sci U S A* 2011; 108:11602-7.
- Loirat C, Veyradier A, Girma JP, Ribba AS, Meyer D. Thrombotic thrombocytopenic purpura associated with von Willebrand factor-cleaving protease (ADAMTS13) deficiency in children. *Semin Thromb Hemost* 2006; 32:90-7.
- Tsai HM. The kidney in thrombotic thrombocytopenic purpura. *Minerva Med* 2007; 98:731-47.
- Rock GA, Shumak KH, Buskard NA, et al. Comparison of plasma exchange with plasma infusion in the treatment of thrombotic thrombocytopenic purpura. Canadian Apheresis Study Group. *N Engl J Med* 1991; 325:393-7.
- Scully M, Yarranton H, Liesner R, et al. Regional UK TTP registry: correlation with laboratory ADAMTS 13 analysis and clinical features. *Br J Haematol* 2008; 142:819-26.
- van de Born BJ, van der Hoeven NV, Groot E, et al. Association between thrombotic microangiopathy and reduced ADAMTS13 activity in malignant hypertension. *Hypertension* 2008; 51:862-6.
- Kremer Hovinga JA, Lämmle B. Role of ADAMTS13 in the pathogenesis, diagnosis, and treatment of thrombotic thrombocytopenic purpura. *Hematology Am Soc Hematol Educ Program* 2012; 2012:610-6.
- Shah N, Rutherford C, Matevosyan K, Shen YM, Sarode R. Role of ADAMTS13 in the management of thrombotic microangiopathies including thrombotic thrombocytopenic purpura (TTP). *Br J Haematol* 2013; 163:514-9.
- Scully M, Knöbl P, Kentouche K, et al. Recombinant ADAMTS-13: first-in-human pharmacokinetics and safety in congenital thrombotic thrombocytopenic purpura. *Blood* 2017; 130:2055-63.
- Cataland SR, Wu HM. How I treat: the clinical differentiation and initial treatment of adult patients with atypical hemolytic uremic syndrome. *Blood* 2014; 123:2478-84.
- Page EE, Kremer Hovinga JA, Terrell DR, Vesely SK, George JN. Thrombotic thrombocytopenic purpura: diagnostic criteria, clinical features, and long-term outcomes from 1995 through 2015. *Blood Adv* 2017; 1:590-600.
- Veyradier A, Obert B, Houllier A, Meyer D, Girma JP. Specific von Willebrand factor-cleaving protease in thrombotic microangiopathies: a study of 111 cases. *Blood* 2001; 98:1765-72.
- Uemura M, Fujimura Y, Matsumoto M, et al. Comprehensive analysis of ADAMTS13 in patients with liver cirrhosis. *Thromb Haemost* 2008; 99:1019-29.

14. Vesely SK, George JN, Lämmle B, et al. ADAMTS13 activity in thrombotic thrombocytopenic purpura – hemolytic uremic syndrome: relation to presenting features and clinical outcomes in a prospective cohort of 142 patients. *Blood* 2003; 102:60-8.
15. Zheng XL, Kaufman RM, Goodnough LT, et al. Effect of plasma exchange on plasma ADAMTS13 metalloprotease activity, inhibitor level, and clinical outcome in patients with idiopathic and nonidiopathic thrombotic thrombocytopenic purpura. *Blood* 2004; 103:4043-9.
16. Mori Y, Wada H, Gabazza EC, et al. Predicting response to plasma exchange in patients with thrombotic thrombocytopenic purpura with measurement of vWF-cleaving protease activity. *Transfusion* 2002; 42:572-80.
17. Franchini M, Mannucci PM. Advantages and limits of ADAMTS13 testing in thrombotic thrombocytopenic purpura. *Blood Transfus* 2008; 6:127-35.
18. Ferrari S, Scheifflinger F, Rieger M, et al; French Clinical and Biological Network on Adult Thrombotic Microangiopathies. Prognostic value of anti-ADAMTS13 antibody features (Ig isotype, titer, and inhibitory effect) in a cohort of 35 adult French patients undergoing a first episode of thrombotic microangiopathy with undetectable ADAMTS13 activity. *Blood* 2007; 109:2815-22.
19. Kremer Hovinga JA, Vesely SK, Terrell DR, Lämmle B, George JN. Survival and relapse in patients with thrombotic thrombocytopenic purpura. *Blood* 2010; 115:1500-11.
20. Scully M. Inhibitory anti-ADAMTS 13 antibodies: measurement and clinical application. *Blood Rev* 2010; 24:11-6.
21. Peyvandi F, Lavoretano S, Palla R, et al. ADAMTS13 and anti-ADAMTS13 antibodies as markers for recurrence of acquired thrombotic thrombocytopenic purpura during remission. *Haematologica* 2008; 93:232-9.
22. Bettoni G, Palla R, Valsecchi C, et al. ADAMTS-13 activity and autoantibodies classes and subclasses as prognostic predictors in acquired thrombotic thrombocytopenic purpura. *J Thromb Haemost* 2012; 10:1556-65.
23. Page EE, Kremer Hovinga JA, Terrell DR, Vesely SK, George JN. Rituximab reduces risk for relapse in patients with thrombotic thrombocytopenic purpura. *Blood* 2016; 127:3092-4.
24. Scully M. Rituximab in the treatment of TTP. *Hematology* 2012; 17 Suppl 1:S22-4.
25. Jang MJ, Kim NK, Chong SY, et al. Frequency of Pro475Ser polymorphism of ADAMTS13 gene and its association with ADAMTS-13 activity in the Korean population. *Yonsei Med J* 2008; 49:405-8.
26. Miyata T, Kokame K, Banno F. Measurement of ADAMTS13 activity and inhibitors. *Curr Opin Hematol* 2005; 12:384-9.
27. Terrell DR, Motto DG, Kremer Hovinga JA, et al. Blood group O and black race are independent risk factors for thrombotic thrombocytopenic purpura associated with severe ADAMTS13 deficiency. *Transfusion* 2011; 51:2237-43.
28. Mayo Clinic Laboratories. Test ID: ADAM13. ADAMTS13 Activity and Inhibitor profile, plasma. Available at: <https://www.mayocliniclabs.com/test-catalog/Overview/61212>. Accessed July 25, 2018.
29. Peyvandi F, Palla R, Lotta LA, et al. ADAMTS-13 assays in thrombotic thrombocytopenic purpura. *J Thromb Haemost* 2009; 8:631-40.
30. Miyata T, Kokame K, Matsumoto M, Fujimura Y. ADAMTS13 activity and genetic mutations in Japan. *Hamostaseologie* 2013; 33:131-7.
31. Feys HB, Liu F, Dong N, et al. ADAMTS-13 plasma level determination uncovers antigen absence in acquired thrombotic thrombocytopenic purpura and ethnic differences. *J Thromb Haemost* 2006; 4:955-62.
32. Cincinnati Children's Clinical Laboratories. Thrombotic microangiopathy (aHUS and TTP) test requisition. Available at: <https://www.cincinnatichildrens.org/-/media/cincinnati%20childrens/home/service/t/thrombotic-microangiopathy/requisitions/tma%20test%20requisition.pdf?la=en>. Accessed July 25, 2018.
33. Cleveland Clinic Laboratories. ADAMTS13 activity and inhibitor assays. Available at: <http://portals.clevelandclinic.org/portals/66/pdf/technical-briefs/adamts13-activity-and-inhibitor-assays.pdf>. Accessed July 25, 2018.



#### About the First Author

Dr Allison Ching Yee Tso is a Senior Consultant Haematologist and Adjunct Assistant Professor at Nanyang Technological University's Lee Kong Chian School of Medicine, and Senior Clinical Lecturer at NUS Yong Loo Lin School of Medicine, Singapore. Dr Tso graduated from the University of Oxford and was trained in London. She is a Fellow of the Royal Colleges of Physicians of London and Edinburgh and a Fellow of the Royal College of Pathologists (United Kingdom). Her research interests include thrombotic microangiopathies, myeloma, haemophagocytic lymphohistiocytosis and haemoglobinopathies.